

CAUTION

KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY



Growth and Finishing Implants for Steers

Each implant (7 pellets) contains: **140 mg trenbolone acetate**
28 mg oestradiol

This product contains a palpable marker.
This product must only be supplied in accordance with Division 4.2 of
the Agvet Code Regulations (Supply of Hormonal Growth Promotants).

**FOR INCREASED RATE OF WEIGHT GAIN &
IMPROVED FEED EFFICIENCY IN STEERS**

Contents: 20 Implants

APVMA Approval No.: 65449/124994

Store between 2 °C and 8 °C (refrigerate, do not freeze).
Protect from light. Store in original container.

For expiry date and batch number see reverse

TEAR HERE

DIRECTIONS FOR USE

To be used **ONLY** in accordance with the instructions on this label unless authorised under
appropriate legislation.

RESTRAINTS

DO NOT USE in any animal other than steers producing beef for human consumption.

DO NOT USE more than one implant for any type concurrently per steer.

DO NOT implant sooner than 70 days after the last implant.

DO NOT implant this product in any site other than beneath the skin in the middle third of the
back of the ear.

DO NOT USE this product on HGP-free accredited cattle. It is a breach of Department of
Agriculture accreditation conditions to use this product on HGP-free accredited cattle or to
possess this product on any HGP-free accredited property (except for feedlots with flexible
accreditation). Treated animals must be identified as having received a hormonal implant by
use of an approved ear punch.

PRECAUTIONS

Implanting cattle under 3 months of age for grass fed beef, or 6 months of age for lot fed beef is NOT
recommended. Implanting cattle 6 weeks of age is NOT recommended.

Implanting cattle intended for breeding is NOT recommended as they may experience decreased fertility.
Growth responses may vary widely for nutritional or other reasons. Improved performance may be noted
only when conditions are favourable for good growth.

Rectal and preputial prolapse, increased bulling activity, high tailheads, sunken loins, ventral oedema
and udder development may occur as side effects of treatment.

To help avoid abscess formation at the site of implant, the surface of the ear should be thoroughly
cleaned and disinfected before implantation.

TEAR HERE

Dosage and Administration:

READ THE ENCLOSED LEAFLET BEFORE USING THIS PRODUCT

Approved implantation technique is fully described in the enclosed leaflet.

Cattle implanted with this product must be individually identified with an approved ear punch mark.

Implant complete contents of a single implant (7 pellets plus marker) per animal.

Implant side is beneath the skin on the back side of the middle one-third of the ear.

WITHHOLDING PERIOD: MEAT: Zero (0) days

TRADE ADVICE:

TRADE ADVICE-EXPORT SLAUGHTER INTERVAL (ESI): An ESI has not established for this product. Note-observing the
meat withholding period may not be sufficient to mitigate potential risks to export trade. Trade advice should be sought from
Elanco Australasia Pty Ltd on 1800 226 324 before using this product.

First Aid Instructions: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 1126

Disposal: **DO NOT** dispose of unused implants in any way other than surrender to an approved retailer or
wholesaler. Dispose of empty cartridges by wrapping with paper and putting in garbage. Discarded needles
should immediately be placed in a designated 'sharps' container.

Manufactured in USA for
Elanco Australasia Pty Ltd
A.B.N. 39 000 233 992
7 Eden Park Drive, Macquarie Park, NSW 2113

May cause cancer.
May damage fertility or the unborn child.
May cause harm to breast-fed children.
(Exposure risk applies if cartridge belt is damaged)

FL102703X

Expiry:

[B]