

Right for cattle. Right by you.

ElancoTM

2024 Elanco Fall Run Cattle Portfolio Offer for Producers



August 1-October 31, 2024

Earn up to a 22% rebate when you spend across multiple product categories with Elanco.*

Buy from two categories



Receive a 12% rebate.

Buy from three categories



Receive a 16% rebate.

Buy from four categories



Receive a 22% rebate.

Producers must purchase a minimum amount per category to earn a rebate:

| Category | Brands | Minimum Spend |
|----------------------------|---|---------------|
| Anti-Infectives | Baytril® 100 (enrofloxacin) Increxxa® (tulathromycin injection) Micotil® (tilmicosin injection) Loncor® (florfenicol) | \$5,000 |
| Vaccines & Immunostimulant | NUPLURA® Titanium® Zelnate® DNA Immunostimulant | \$2,500 |
| Implants | Component E-C Component TE-G Compudose® Encore® All Component with Tylan SKUs (progesterone and estradiol benzoate and tylosin tartrate implants) & (trenbolone acetate and estradiol and tylosin tartrate implants) | \$2,500 |
| Cydectin & Fly Control | Cydectin® (moxidectin) Injectable Cydectin® (moxidectin) Pour-On Cydectin® (moxidectin) Oral Sheep Drench Clean-Up II™ Pour-On Insecticide with IGR Corathon® CyLence® Pour-On Insecticide CyLence Ultra® Insecticide Cattle Ear Tag Patriot® Insecticide Cattle Ear Tag Permethrin® II Permethrin® II CDS Pour-On Rabon® 50 WP Insecticide | \$2,500 |



Earn an additional 3% rebate on Cydectin Injectable when you spend \$1000 on Cydectin Injectable.



ElancoCattleRebates.com

Program Terms & Conditions

Ways to claim your rebate:

1. Online claims: Scan the QR code to be directed to the rebate claim forms online, or visit ElancoCattleRebates.com.
2. Mail-in claims: Complete a mail-in claim form and submit with proof of purchase (invoice) via mail.
3. Supplier Reported Claims (EDI): Qualifying purchases billed through an EDI-reporting distributor be aggregated at the end of the program. EDI-reporting distributor must report EDI sales to Elanco. Qualifying customers will automatically receive a rebate check in the mail.

Program Terms

- Only producers are eligible for this program. Producers enrolled in any special Elanco pricing programs are not eligible for this rebate offer. Producers participating in the 2024 Elanco Cattle Fall Bonus Brand Savings for Producers are not eligible for this rebate offer.
- Purchases are cumulative August 1- October 31, 2024. Eligible purchases must be invoiced by October 31, 2024.
- Minimum rebate check amount is \$50.00. Allow at minimum 6 weeks for processing following program conclusion.

- Eligible products for the offer include: Baytril 100 (enrofloxacin), Increxxa (tulathromycin injection), Loncor 300 (florfenicol), Micotil (tilmicosin injection), Zelnate, Cydectin Injectable, Cydectin Pour On, Cydectin Drench, Titanium, Nuplura PH, Component E-C, Component E-C with Tylan, Component TE-G, Component TE-G with Tylan, Component TE-IH with Tylan, Component TE-IS with Tylan, Component TE-200 with Tylan Compudose, Encore, Clean up™ II Pour-on Insecticide with IGR, Corathon®, CyLence® Pour-On Insecticide, CyLence Ultra® Insecticide Cattle Ear Tag, Patriot® Insecticide Cattle Ear Tag, Permethrin® CDS Pour-on Insecticide, Permethrin® II, Rabon® 50 WP Insecticide. All sizes of each SKU are eligible. Offers valid while supplies last.
- Products that are on backorder and qualify for the rebate offer must ship during the program term to receive the rebate. Products that are on backorder and do not ship during the program term will not qualify for any rebate in this program. All product returns will be applied toward the subsequent program payout period.
- Producers who submit coupons via mail or online will not be eligible for EDI reported claims.
- Limit 10 mail or online claim submissions per household. Multiple proofs of purchase can be used per submission. One rebate will be calculated per submission. Qualifying rebates will be calculated for all invoices included in a submission and will not be calculated across multiple submissions.
- Mail and online rebate claim form and invoice(s) must be postmarked or submitted by November 30, 2024.
- Elanco reserves the right to vary the terms and conditions of this program or to cancel this program at any time upon notice through the website Elanco.com.
- For more information regarding this rebate program and eligibility, contact your Elanco Animal Health representative or call 800-364-2014.

| Categories | Category Minimum Spend | Product | Amount Purchased | | Rebate Percent | | Total \$ |
|---|------------------------|---|------------------|---|----------------|---|-----------|
| Anti-infectives | \$5,000 | Baytril 100 (enrofloxacin) | | x | | = | \$ |
| | | Increxxa (tulathromycin injection) | | x | | = | \$ |
| | | Micotil (tilmicosin injection) | | x | | = | \$ |
| | | Loncor 300 (florfenicol) | | x | | = | \$ |
| Vaccines & Immunostimulant | \$2,500 | Nuplura PH | | x | | = | \$ |
| | | Titanium | | x | | = | \$ |
| | | Zelnate | | x | | = | \$ |
| Implants | \$2,500 | Component E-C | | x | | = | \$ |
| | | Component TE-G | | x | | = | \$ |
| | | All Component with Tylan SKUs | | x | | = | \$ |
| | | Compudose | | x | | = | 4 |
| | | Encore | | x | | = | \$ |
| Cydectin & Fly Control | \$2,500 | Cydectin Injectable | | x | | = | \$ |
| | | Cydectin Pour-On | | x | | = | \$ |
| | | Cydectin Drench | | x | | = | \$ |
| | | Clean Up II™ Pour-on Insecticide with IGR | | x | | = | \$ |
| | | Corathon® | | x | | = | \$ |
| | | CyLence® Pour-On Insecticide | | x | | = | \$ |
| | | CyLence® Ultra Insecticide Cattle Ear Tag | | x | | = | \$ |
| | | Patriot® Insecticide Cattle Ear Tag | | x | | = | \$ |
| | | Permethrin® CDS Pour-on Insecticide | | x | | = | \$ |
| | | Permethrin® II | | x | | = | \$ |
| Rabon® 50 WP Insecticide | | x | | = | \$ | | |
| Earn an additional 3% rebate on Cydectin Injectable when you spend \$1000 on Cydectin Injectable. | | | | x | 3% | = | \$ |
| TOTAL Rebate Request: | | | | | | | \$ |

Submit rebate claim form and invoice(s)
POSTMARKED NO LATER THAN June 30, 2024 to:

Invoice copy(ies) or invoice statement(s) must accompany this form to obtain rebate.
Invoices or statements must clearly show:

Elanco Cydectin & Insecticides
Spring 2024 Rebate
P.O. Box 1080, Dept. R12992
Grand Rapids, MN 55745-1080

1. Product name(s), product size(s) and date(s) purchased.
2. Name of company where product(s) was/were purchased and price paid.
3. No limit to quantity of rebate claims.

Producer Name _____

Physical Address _____

Mailing Address _____

City _____ State _____ Zip Code _____

Phone Number _____

Email Address _____

I would like to receive future communications from Elanco about farm animal products.

For all products: The label contains complete use information, including cautions and warnings. Always read, understand and follow the label and use directions.

Keep Cydectin out of reach of children.

For all Component with Tylan implants:

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

IMPORTANT SAFETY INFORMATION FOR MICOTIL Before using this product, it is important to read the entire product insert, including the boxed human warning.

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian. Not for human use. Injection of this drug in humans has been associated with fatalities. Keep out of reach of children. Administer only with a tube-fed safety syringe. Do not use in automatically powered syringes, single-use syringes, or other delivery devices. Exercise extreme caution to avoid accidental self-injection. In case of human injection, consult a physician immediately and apply ice or cold pack to injection site while avoiding direct contact with the skin. Avoid contact with eyes. Always use proper drug handling procedures to avoid accidental self-injection. Consult your veterinarian on the safe handling and use of all injectable products prior to administration. For use in cattle or sheep only. Inject subcutaneously. Injection of this antibiotic has been shown to be fatal in swine and non-human primates, and may be fatal in horses and goats. Do not use in lambs less than 15 kg body weight. Do not use in female dairy cattle 20 months of age or older. Use in lactating dairy cattle or sheep may cause milk residues. The following adverse reactions have been reported: in cattle: injection site swelling and inflammation, lameness, collapse, anaphylaxis/anaphylactoid reactions, decreased food and water consumption, and death; in sheep: dyspnea and death. Micotil has a pre-slaughter withdrawal time of 42 days

Zelnate is based on technology developed by Juvaris BioTherapeutics and is patent protected. Animal health applications are being developed exclusively under the rights of Elanco and are protected by patents.

Baytril, Loncor and QuickBayt are sold by Elanco or its affiliates and are not Bayer products. The Baytril, Loncor and QuickBayt trademarks are owned by Bayer and used under license.

Clean-Up, Component, Compudose, Corathon, Cydectin, CyLence, Encore, Increxxa, Micotil, Nuplura PH+, Patriot, Permethrin, Rabon, Titanium, Tylan, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

FULL PRESCRIPTION INFORMATION FOR USE IN CATTLE ONLY. SEE PRODUCT INSERT FOR COMPLETE DOSING AND ADMINISTRATION INFORMATION.

Elanco™

Micotil™ 300™

250 mL

(tilmicosin injection)

300 mg tilmicosin, USP as tilmicosin phosphate per mL
For Subcutaneous Use in Cattle Only

Administer only with a tube-fed safety syringe. Do not use in automatically powered syringes, single-use syringes, or other delivery devices. Contact Elanco at 1-800-428-4441, or your distributor, for a tube-fed safety syringe for use with this product.

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian. Before using Micotil, please consult the product insert, a summary of which follows:

Indications: For the treatment of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni*. For the control of respiratory disease in cattle at high risk of developing BRD associated with *Mannheimia haemolytica*.

Approved by FDA under NADA # 140-929

Micotil must be used with the quick-fit connector made specifically for its use. Contact Elanco or your distributor for this equipment. Read product labeling, including Safe Handling Practices, before use.

Dosage and Administration: Follow instructions for activation of the shroud before first usage. Inject Subcutaneously in Cattle Only. See Safe Handling Practices, Contraindications, and Warnings prior to use. In cattle, administer a single subcutaneous dose of 10 to 20 mg/kg body weight (1 to 2 mL/30 kg or 1.5 to 3 mL per 100 lbs).

Do not inject more than 10 mL per injection site. If no improvement is noted within 48-hours, the diagnosis should be reevaluated. For cattle injection under the skin in the neck is suggested. If not accessible, inject under the skin behind the shoulders and over the ribs.

Note: Swelling at the subcutaneous site of injection may be observed. See product insert for complete dosing and administration information.

CONTRAINDICATIONS: Do not use in automatically powered syringes, single-use syringes, or other delivery devices not specified in the labeling. Do not administer intravenously to cattle. Intravenous injection in cattle will be fatal. Do not administer to animals other than cattle. Injection of tilmicosin has been shown to be fatal in swine and non-human primates. Death following exposure to tilmicosin injection has been reported to FDA/CVM in goats, rabbits, pheasants, pigs, dogs, deer, cats, alpacas, and horses.

Residue Warnings: Animals intended for human consumption must not be slaughtered within 42 days of the last treatment. Not for use in lactating dairy cattle 20 months of age or older. Use of tilmicosin in this class of cattle may cause milk residues.

Precautions: The effects of tilmicosin on bovine reproductive performance, pregnancy and lactation have not been determined. Intramuscular injection will cause a local reaction which may result in trim loss of edible tissue at slaughter.

Storage Conditions: Store at or below 86 °F (30 °C). Protect from direct sunlight. Use within 84 days of first puncture. Date of first puncture:

To report adverse effects, access medical information, or obtain additional product information, call 1-800-428-4441.

HUMAN WARNINGS: Not for human use. Injection of this drug in humans has been associated with fatalities. Keep out of reach of children. Administer only with a tube-fed safety syringe. Do not use in automatically powered syringes, single-use syringes, or other delivery devices. Exercise extreme caution to avoid accidental self-injection. In case of human injection, consult a physician immediately and apply ice or cold pack to injection site while avoiding direct contact with the skin. Emergency medical telephone numbers are 1-800-722-0987 or 1-800-428-4441. Avoid contact with skin, eyes or mucous membranes.

NOTE TO THE PHYSICIAN: The cardiovascular system is the target of toxicity and should be monitored closely. Cardiovascular toxicity may be due to calcium channel blockade. In dogs, administration of intravenous calcium offset Micotil-induced tachycardia and negative inotropy (decreased contractility). Dobutamine partially offset the negative inotropic effects induced by Micotil in dogs. β-adrenergic antagonists, such as propranolol, exacerbated the negative inotropy of Micotil in dogs. Epinephrine potentiated lethality of Micotil in pigs. This antibiotic persists in tissues for several days.

Adverse Reactions: The following adverse reactions have been reported post-approval: In cattle: injection site swelling and inflammation, lameness, collapse, anaphylaxis/anaphylactoid reactions, decreased food and water consumption, and death. For additional information about reporting adverse drug experiences for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

Effectiveness: In a multi-location field study, 1508 calves with naturally occurring BRD were treated with Micotil. Responses to treatment were compared to saline-treated controls. A cure was defined as a calf with normal attitude and activity, normal respiration, and a rectal temperature of <104°F on Day 13. The cure rate was significantly higher (P=0.004) in Micotil-treated calves (63.1%) compared to saline-treated calves (29.2%). During the treatment phase of the study, there were 10 BRD-related deaths in the Micotil-treated calves compared to 47 in the saline-treated calves.

How Supplied: Micotil (tilmicosin injection) is supplied in 250 mL multi-dose amber glass bottles in a non-removable polymer protector.

Manufactured by:
Elanco US, Inc. Greenfield, IN 46140, USA

Revised: 09/2021

Micotil, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

© 2023 Elanco or its affiliates. PM-US-23-1619

Instructions for Activation of the Shroud

Before first usage activate the shroud-vial-system as shown in the pictures. Administer only with a tube-fed safety syringe. Do not use in automatically powered syringes, single-use syringes, or other delivery devices. This product must be used with the quick-fit connector made specifically for use with Micotil (tilmicosin injection) that attaches to the shroud fitting. To obtain a tube-fed safety syringe and quick-fit connector, contact Elanco at 1-800-428-4441, or your distributor

Step 1.


Twist the two tamper-evident tabs to break them off the Shroud Base.

Step 2.

Rotate the Shroud Top through a quarter-turn clockwise. The spike will pierce the vial closure, and the Shroud Top will lock into its final position by an audible "click".

Step 3.

The correct final position can be confirmed by the alignment of the 2 arrows as shown in the picture.




Step 4.

Remove the flexible cap from the fluid connection. Attach the quick-fit connector to tubing if not already attached. Push the quick-fit connector downwards onto the shroud fitting until it clicks into place.

Step 5.

Invert the Micotil Shroud, then prime the tube-fed safety syringe following manufacturer's instructions.

Return shroud to upright position after finishing operation. Leave tubing attached to tube-fed safety syringe and quick-fit connector until dosing equipment has been removed from the shroud. Remove dosing equipment by pushing the trigger as shown in the picture, then disconnecting the quick-fit connector from the shroud.



Micotil should not be stored in dosing equipment. Dosing equipment should be disconnected from the shroud after each use. Store product upright. The dosing equipment should be cleaned according to the manufacturer's instructions. Avoid contact with skin, eyes, or mucous membranes.

1. WHAT ARE THE POSSIBLE EFFECTS OF ACCIDENTAL HUMAN INJECTION?

Human injections of Micotil have been associated with fatalities. Clinical signs from human exposure include od taste in the mouth, nausea, headache, dizziness, rapid heart rate, chest pain, anxiety, or lightheadedness. Local reactions such as injection site pain, bleeding, swelling or inflammation have been reported.

2. WHAT SHOULD I DO IN THE CASE OF ACCIDENTAL HUMAN INJECTION?

- Immediately seek medical attention.
- Apply ice or cold pack to injection site, while avoiding direct contact with the skin, and transport immediately to a hospital.
- Call 1-800-722-0987 or 1-800-428-4441 for further emergency information.

3. WHAT SHOULD MY PHYSICIAN KNOW IN THE CASE OF ACCIDENTAL HUMAN INJECTION?


- The cardiovascular system is the target of toxicity and should be monitored closely.
- Cardiovascular toxicity may be due to calcium channel blockade.
- Intravenous calcium administration reversed the cardiovascular effects of Micotil in dogs and may provide benefit in patients exhibiting low blood pressure (hypotension) or rapid heart rate (tachycardia).
- Dobutamine improved some of the cardiac function in dogs given Micotil.
- Epinephrine increased the toxicity of Micotil in pigs, resulting in death.
- Propranolol (a beta-adrenergic antagonist) further decreased cardiac function in dogs given Micotil.
- The active ingredient in Micotil is tilmicosin phosphate and persists in tissue for several days.
- Call 1-800-722-0987 or 1-800-428-4441 for further emergency information.

4. WHAT ARE THE PROPER WAYS TO HANDLE AND STORE MICOTIL?

- Store at or below 86°F (30°C), out of direct sunlight, in a safe location, not easily accessible to the general public. Use within 84 days of first puncture. Store upright between product dispensing. Disconnect and clean dosing equipment for storing as per manufacturer's instructions.
- Avoid contact with skin, eyes, or mucous membranes.
- Read, understand, and follow all label use directions.
- Wash hands thoroughly with soap and water after handling.

5. WHAT ARE THE PROPER METHODS FOR ADMINISTERING MICOTIL?

- Properly restrain animals prior to administration.
- Work in a team, or if alone, advise someone of your location and how long you plan to be there.
- For subcutaneous use. Administer only with a tube-fed safety syringe. Do not use in automatically powered syringes, single-use syringes, or other delivery devices. Contact Elanco at 1-800-428-4441, or your distributor, for a tube-fed safety syringe for use with this product.
- Use a 1/2-inch to 5/8-inch, 18- to 16-gauge needle.
- With a single hand on the safety syringe insert the needle subcutaneously, at a top-down angle, while avoiding penetration of underlying muscle.
- For cattle, injection under the skin in the neck is suggested. If not accessible, inject under the skin behind the shoulders and over the ribs.
- In cattle, administer a single subcutaneous dose of 1.5 to 3.0 mL of Micotil (tilmicosin injection) per 100 lbs of body weight, in either of the two areas noted in the adjacent drawing.
- For beef cattle, Beef Quality Assurance recommends injection site 1, unless this site is inaccessible or places the operator in a potentially dangerous situation.
- Wash hands thoroughly with soap and water after administration.
- Do not administer intravenously (IV) as IV administration will be fatal.
- Intramuscular injection will cause a local reaction, which may result in trim loss.
- Do not inject more than 10 mL per injection site.
- Do not use in lambs less than 15 kg body weight.



6. WHAT ARE SAFE WAYS TO REMOVE AND CHANGE NEEDLES?

- Always follow the manufacturer's instruction of how to safely remove and change needles from the safety syringe.
- Plan for the safe handling and disposal of needles before use.
- Keep the needle capped until ready to use.
- Avoid recapping a used needle.
- To safely remove used needles, use tools appropriate for the specific type of safety syringe. Do not remove a used needle with your fingers.
- Dispose used needles in an appropriate sharps disposal container. Do not overfill sharps containers and do not put your fingers into a sharps container.
- Never place loose needles in household or public trash cans.

**FULL PRESCRIBING INFORMATION FOR USE IN CATTLE ONLY
SEE PRODUCT INSERT FOR COMPLETE DOSING AND
ADMINISTRATION INFORMATION**

**Elanco™
Increxxa™
(tulathromycin injection)**


Injectable Solution

Antibiotic
100 mg of tulathromycin/mL

For use in beef cattle (including suckling calves), non-lactating dairy cattle (including dairy calves) and veal calves. Not for use in female dairy cattle 20 months of age or older.
CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION

Increxxa Injectable Solution is a ready-to-use sterile parenteral preparation containing tulathromycin, a semi-synthetic macrolide antibiotic of the subclass triamides. Each mL of Increxxa contains 100 mg of tulathromycin, 500 mg propylene glycol, 19.2 mg citric acid and 5 mg monothiolglycerol. Sodium hydroxide or hydrochloric acid may be added to adjust pH. Increxxa consists of an equilibrated mixture of two isomeric forms of tulathromycin in a 9:1 ratio. Structures of the isomers are shown below.



The chemical names of the isomers are (2R,3S,4R,5R,6R,10R,11R,12S,13S,14R)-13-[[2,6-dideoxy-3-C-methyl-3-O-methyl-4-C-(propylamino) methyl]-a-L-ribo-hexopyranosyl]-2-ethyl-3,4,10-trihydroxy-3,5,8,10,12,14-hexamethyl-11-[[3,4,6-trideoxy-3-(dimethylamino)-β-D-xyllo-hexopyranosyl]-oxy]-1-oxa-6-azacyclotridecan-15-one and (2R,3R,6R,8R,9R,10S,11S,12R)-11-[[2,6-dideoxy-3-C-methyl-3-O-methyl-4-C-(propylamino)methyl]-a-L-ribo-hexopyranosyl]-2-[(1R,2R)-1,2-dihydroxy-1-methylbutyl]-8-hydroxy-3,6,8,10,12-pentamethyl-9-[[3,4,6-trideoxy-3-(dimethylamino)-β-D-xyllo-hexopyranosyl]-oxy]-1-oxa-4-azacyclotridecan-13-one, respectively.

INDICATIONS

Beef and Non-Lactating Dairy Cattle

BRD – Increxxa Injectable Solution is indicated for the treatment of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida*, *Histophilus somni*, and *Mycoplasma bovis*; and for the control of respiratory disease in cattle at high risk of developing BRD associated with *Mannheimia haemolytica*, *Pasteurella multocida*, *Histophilus somni*, and *Mycoplasma bovis*.

IBK – Increxxa Injectable Solution is indicated for the treatment of infectious bovine keratoconjunctivitis (IBK) associated with *Moraxella bovis*.

Foot Rot – Increxxa Injectable Solution is indicated for the treatment of bovine foot rot (interdigital necrobacillosis) associated with *Fusobacterium necrophorum* and *Porphyromonas levis*.

Suckling Calves, Dairy Calves, and Veal Calves

BRD – Increxxa Injectable Solution is indicated for the treatment of BRD associated with *M. haemolytica*, *P. multocida*, *H. somni*, and *M. bovis*.

DOSEAGE AND ADMINISTRATION

Cattle

Inject subcutaneously as a single dose in the neck at a dosage of 2.5 mg/kg (1.1 mL/100 lb) body weight (BW). Do not inject more than 10 mL per injection site.

Table 1. Increxxa Cattle Dosing Guide

| Animal Weight (Pounds) | Dose Volume (mL) |
|------------------------|------------------|
| 100 | 1.1 |
| 200 | 2.3 |
| 300 | 3.4 |
| 400 | 4.5 |
| 500 | 5.7 |
| 600 | 6.8 |
| 700 | 8.0 |
| 800 | 9.1 |
| 900 | 10.2 |
| 1000 | 11.4 |

CONTRAINDICATIONS

The use of Increxxa Injectable Solution is contraindicated in animals previously found to be hypersensitive to the drug.

WARNINGS

FOR USE IN ANIMALS ONLY.

NOT FOR HUMAN USE.

KEEP OUT OF REACH OF CHILDREN.

NOT FOR USE IN CHICKENS OR TURKEYS.

RESIDUE WARNINGS

Cattle

Cattle intended for human consumption must not be slaughtered within 18 days from the last treatment. This drug is not approved for use in female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows.

PRECAUTIONS

Cattle

The effects of Increxxa on bovine reproductive performance, pregnancy, and lactation have not been determined. Subcutaneous injection can cause a transient local tissue reaction that may result in trim loss of edible tissue at slaughter.

ADVERSE REACTIONS

Cattle

In one BRD field study, two calves treated with tulathromycin injection at 2.5 mg/kg BW exhibited transient hypersalivation. One of these calves also exhibited transient dyspnea, which may have been related to pneumonia.

POST APPROVAL EXPERIENCE

The following adverse events are based on post approval adverse drug experience reporting. Not all adverse events are reported to the FDA CVM. It is not always possible to reliably estimate the adverse event frequency or establish a causal relationship to product exposure using these data. The following adverse events are listed in decreasing order of reporting frequency in cattle: Injection site reactions and anaphylaxis/anaphylactoid reactions. For additional information about reporting adverse drug experiences for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimal>.

CLINICAL PHARMACOLOGY

At physiological pH, tulathromycin (a weak base) is approximately 50 times more soluble in hydrophilic than hydrophobic media. This solubility profile is consistent with the extracellular pathogen activity typically associated with the macrolides. 1. Markedly higher tulathromycin concentrations are observed in the lungs as compared to the plasma. The extent to which lung concentrations represent free (active) drug was not examined. Therefore, the clinical relevance of these elevated lung concentrations is undetermined. Although the relationship between tulathromycin and the characteristics of its antimicrobial e-acts has not been characterized, as a class, macrolides tend to be primarily bacteriostatic, but may be bactericidal against some pathogens. 2. They also tend to exhibit concentration independent killing; the rate of bacterial eradication does not change once serum drug concentrations reach 2 to 3 times the minimum inhibitory concentration (MIC) of the targeted pathogen. Under these conditions, the time that serum concentrations remain above the MIC becomes the major determinant of antimicrobial activity. Macrolides also exhibit a post-antibiotic e-act (PAE), the duration of which tends to be both drug and pathogen dependent. In general, by increasing the macrolide concentration and the exposure time, the PAE will increase to some maximal duration. Of the two variables, concentration and exposure time, drug concentration tends to be the most powerful determinant of the duration of PAE. Tulathromycin is eliminated from the body primarily unchanged via biliary excretion.

¹ Carbon, C. 1998. *Pharmacodynamics of Macrolides, Azalides, and Streptogramins: Effect on Extracellular Pathogens*. Clin. Infect. Dis., 27:28-32.

² Nightingale, C.J. 1997. *Pharmacokinetics and Pharmacodynamics of Newer Macrolides*. Pediatr. Infect. Dis. J., 16:438-443.

Cattle

Following subcutaneous administration into the neck of feeder calves at a dosage of 2.5 mg/kg BW, tulathromycin is rapidly and nearly completely absorbed. Peak plasma concentrations generally occur within 15 minutes after dosing and product relative bioavailability exceeds 90%. Total systemic clearance is approximately 170 mL/hr/kg. Tulathromycin distributes extensively into body tissues, as evidenced by volume of distribution values of approximately 11 L/kg in healthy ruminating calves. ³ This extensive volume of distribution is largely responsible for the long elimination half-life of this compound [approximately 2.75 days in the plasma (based on quantifiable terminal plasma drug concentrations) versus 8.75 days for total lung concentrations (based on data from healthy animals)]. Linear pharmacokinetics are observed with subcutaneous doses ranging from 1.27 mg/kg BW to 5.0 mg/kg BW. No pharmacokinetic differences are observed in castrated male versus female calves.

³ Clearance and volume estimates are based on intersubject comparisons of 2.5 mg/kg BW administered by either subcutaneous or intravenous injection.

MICROBIOLOGY

Cattle

Tulathromycin has demonstrated in vitro activity against *Mannheimia haemolytica*, *Pasteurella multocida*, *Histophilus somni*, and *Mycoplasma bovis*, four pathogens associated with BRD; against *Moraxella bovis* associated with IBK; and against *Fusobacterium necrophorum* and *Porphyromonas levis* associated with bovine foot rot. The MICs of tulathromycin against indicated BRD and IBK pathogens were determined using methods recommended by the Clinical and Laboratory Standards Institute (CLSI, M31-A2). The MICs against foot rot pathogens were also determined using methods recommended by the CLSI (M11-A6). All MIC values were determined using the 9:1 isomer ratio of this compound.

BRD – The MICs of tulathromycin were determined for BRD isolates obtained from calves enrolled in therapeutic and at-risk field studies in the U.S. in 1999. In the therapeutic studies, isolates were obtained from pre-treatment nasopharyngeal swabs from all study calves, and from lung swabs or lung tissue of saline-treated calves that died. In the at-risk studies, isolates were obtained from nasopharyngeal swabs of saline-treated non-responders, and from lung swabs or lung tissue of saline-treated calves that died. The results are shown in Table 3.

IBK – The MICs of tulathromycin were determined for *Moraxella bovis* isolates obtained from calves enrolled in IBK field studies in the U.S. in 2004. Isolates were obtained from pre-treatment conjunctival swabs of calves with clinical signs of IBK enrolled in the tulathromycin injection and saline-treated groups. The results are shown in Table 3.

Foot Rot – The MICs of tulathromycin were determined for *Fusobacterium necrophorum* and *Porphyromonas levis* obtained from cattle enrolled in foot rot field studies in the U.S. and Canada in 2007. Isolates were obtained from pre-treatment interdigital biopsies and swabs of cattle with clinical signs of foot rot enrolled in the tulathromycin injection and saline-treated groups. The results are shown in Table 3.

Table 3. Tulathromycin minimum inhibitory concentration (MIC) values* for indicated pathogens isolated from field studies evaluating BRD and IBK in the U.S. and from foot rot field studies in the U.S. and Canada.

| Indicated pathogen | Date isolated | No. of isolates | MIC ₅₀ ** (µg/mL) | MIC ₉₀ ** (µg/mL) | MIC range (µg/mL) |
|----------------------------------|---------------|-----------------|------------------------------|------------------------------|-------------------|
| <i>Mannheimia haemolytica</i> | 1999 | 642 | 2 | 2 | 0.5 to 64 |
| <i>Pasteurella multocida</i> | 1999 | 221 | 0.5 | 1 | 0.25 to 64 |
| <i>Histophilus somni</i> | 1999 | 36 | 4 | 4 | 1 to 4 |
| <i>Mycoplasma bovis</i> | 1999 | 43 | 0.125 | 1 | ≤ 0.063 to > 64 |
| <i>Moraxella bovis</i> | 2004 | 55 | 0.5 | 0.5 | 0.25 to 1 |
| <i>Fusobacterium necrophorum</i> | 2007 | 116 | 2 | 64 | ≤ 0.25 to > 128 |
| <i>Porphyromonas levis</i> | 2007 | 103 | 8 | 128 | ≤ 0.25 to > 128 |

* The correlation between in vitro susceptibility data and clinical e-ectiveness is unknown.
** The lowest MIC to encompass 50% and 90% of the most susceptible isolates, respectively.

EFFECTIVENESS

Cattle

BRD – In a multi-location field study, 314 calves with naturally occurring BRD were treated with tulathromycin injection. Responses to treatment were compared to saline-treated controls. A cure was defined as a calf with normal attitude/activity, normal respiration, and a rectal temperature of ≤ 104°F on Day 14. The cure rate was significantly higher (P ≤ 0.05) in tulathromycin injection-treated calves (78%) compared to saline-treated calves (24%). There were two BRD-related deaths in the tulathromycin injection-treated calves compared to nine BRD-related deaths in the saline-treated calves. Fifty-two tulathromycin injection-treated calves and 27 saline-treated calves from the multi-location field BRD treatment study had *Mycoplasma bovis* identified in cultures from pre-treatment nasopharyngeal swabs.

Of the 52 tulathromycin injection-treated calves, 37 (71.2%) calves were categorized as cures and 15 (28.8%) calves were categorized as treatment failures. Of the 27 saline-treated calves, 4 (14.8%) calves were categorized as cures and 23 (85.2%) calves were treatment failures.

A Bayesian meta-analysis was conducted to compare the BRD treatment success rate in young calves (calves weighing 250 lbs or less and fed primarily a milk-based diet) treated with tulathromycin injection to the success rate in older calves (calves weighing more than 250 lbs and fed primarily a roughage and grain-based diet) treated with tulathromycin injection. The analysis included data from four BRD treatment effectiveness studies conducted for the approval of tulathromycin injection in the U.S. and nine contemporaneous studies conducted in Europe. The analysis showed that the BRD treatment success rate in young calves was at least as good as the BRD treatment success rate in older calves. As a result, tulathromycin injection is considered effective for the treatment of BRD associated with *M. haemolytica*, *P. multocida*, *H. somni*, and *M. bovis* in suckling calves, dairy calves, and veal calves.

In another multi-location field study with 399 calves at high risk of developing BRD, administration of tulathromycin injection resulted in a significantly reduced incidence of BRD (11%) compared to saline-treated calves (59%). Effectiveness evaluation was based on scored clinical signs of normal attitude/activity, normal respiration, and a rectal temperature of ≤ 104°F on Day 14. There were no BRD-related deaths in the tulathromycin injection-treated calves compared to two BRD-related deaths in the saline-treated calves.

Fifty saline-treated calves classified as non-responders in this study had *Mycoplasma bovis* identified in cultures of post-treatment nasopharyngeal swabs or lung tissue.

Two induced infection model studies were conducted to confirm the e-ectiveness of tulathromycin injection against *Mycoplasma bovis*. A total of 166 calves were inoculated intratracheally with field strains of *Mycoplasma bovis*. When calves became pyrexid and had abnormal respiration scores, they were treated with either tulathromycin injection (2.5 mg/kg BW) subcutaneously or an equivalent volume of saline. Calves were observed for signs of BRD for 14 days post-treatment, then were euthanized and necropsied. In both studies, mean lung lesion percentages were statistically significantly lower in the tulathromycin injection-treated calves compared with saline-treated calves (11.3% vs. 28.9%, P = 0.0001 and 15.0% vs. 30.7%, P < 0.0001).

IBK – Two field studies were conducted evaluating tulathromycin injection for the treatment of IBK associated with *Moraxella bovis* in 200 naturally-infected calves. The primary clinical endpoint of these studies was cure rate, defined as a calf with no clinical signs of IBK and no corneal ulcer, assessed on Days 5, 9, 13, 17, and 21. Time to improvement, defined as the first day on which a calf had no clinical signs of IBK in both eyes, provided that those scores were maintained at the next day of observation, was assessed as a secondary variable. At all time points, in both studies, the cure rate was significantly higher (P < 0.05) for tulathromycin injection-treated calves compared to saline-treated calves. Additionally, time to improvement was significantly less (P < 0.0001) in both studies for tulathromycin injection-treated calves compared to saline-treated calves.

Foot Rot – The effectiveness of tulathromycin injection for the treatment of bovine foot rot was evaluated in 170 cattle in two field studies. Cattle diagnosed with bovine foot rot were enrolled and treated with a single subcutaneous dose of tulathromycin injection (2.5 mg/kg BW) or an equivalent volume of saline. Cattle were clinically evaluated 7 days after treatment for treatment success, which was based on defined decreases in lesion, swelling, and lameness scores. In both studies, the treatment success percentage was statistically significantly higher in tulathromycin injection-treated calves compared with saline-treated calves (60% vs. 8%, P < 0.0001 and 83.3% vs. 50%, P = 0.0088).

ANIMAL SAFETY

Cattle

Safety studies were conducted in feeder calves receiving a single subcutaneous dose of 25 mg/kg BW, or 3 weekly subcutaneous doses of 2.5, 7.5, or 12.5 mg/kg BW. In all groups, transient indications of pain after injection were seen, including head shaking and pawing at the ground. Injection site swelling, discoloration of the subcutaneous tissues at the injection site and corresponding histopathologic changes were seen in animals in all dosage groups. These lesions showed signs of resolving over time. No other drug-related lesions were observed macroscopically or microscopically. An exploratory study was conducted in feeder calves receiving a single subcutaneous dose of 10, 12.5, or 15 mg/kg BW. Macroscopically, no lesions were observed. Microscopically, minimal to mild myocardial degeneration was seen in one of six calves administered 12.5 mg/kg BW and two of six calves administered 15 mg/kg BW.

A safety study was conducted in pre-ruminant calves 13 to 27 days of age receiving 2.5 mg/kg BW or 7.5 mg/kg BW once subcutaneously. With the exception of minimal to mild injection site reactions, no drug-related clinical signs or other lesions were observed macroscopically or microscopically.

STORAGE CONDITIONS

Store below 25°C (77°F), with excursions up to 40°C (104°F).

100 mL: Use within 2 months of first puncture and puncture a maximum of 67 times.

If more than 67 punctures are anticipated, the use of multi-dosing equipment is recommended. When using a draw-off spike or needle with bore diameter larger than 16 gauge, discard any product remaining in the vial immediately after use.

250 mL: Use within 2 months of first puncture and puncture a maximum of 100 times.

If more than 100 punctures are anticipated, the use of multi-dosing equipment is recommended. When using a draw-off spike or needle with bore diameter larger than 16 gauge, discard any product remaining in the vial immediately after use.

HOW SUPPLIED

Increxxa (tulathromycin injection) Injectable Solution is available in the following package sizes:

100 mL vial

250 mL vial

500 mL vial

For product questions, to report adverse reactions, or for a copy of the Safety Data Sheet (SDS), call Elanco Product & Veterinary Support at 1-800-428-4441. For additional information about reporting adverse drug experiences for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimal>. Approved by FDA under ANADA # 200-666 Product of China.

Manufactured for: Elanco US Inc., Greenfield, IN 46140 U.S.A.

Increxxa, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

© 2023 Elanco or its affiliates. PM-US-23-1622





**100 mg/mL Antimicrobial
Injectable Solution**

For Subcutaneous Use In Beef Cattle And Non-Lactating Dairy Cattle
Not For Use In Female Dairy Cattle 20 Months Of Age Or Older
Or In Calves To Be Processed For Veal

CAUTION:

Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian.
Federal (U.S.A.) law prohibits the extra-label use of this drug in food-producing animals.

Before Using Baytril 100, please consult the complete product insert, a summary of which follows:

INDICATIONS:

Cattle - Single-Dose Therapy: Baytril 100 is indicated for the treatment of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida*, *Histophilus somni* and *Mycoplasma bovis* in beef and non-lactating dairy cattle; and for the control of BRD in beef and non-lactating dairy cattle at high risk of developing BRD associated with *M. haemolytica*, *P. multocida*, *H. somni* and *M. bovis*.

Cattle - Multiple-Day Therapy: Baytril 100 is indicated for the treatment of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni* in beef and non-lactating dairy cattle.

DOSAGE AND ADMINISTRATION:

Baytril 100 provides flexible dosages and durations of therapy. Baytril 100 may be administered as a single dose for one day for treatment and control of BRD (cattle), or for multiple days for BRD treatment (cattle). Selection of the appropriate dose and duration of therapy for BRD treatment in cattle should be based on an assessment of the severity of the disease, pathogen susceptibility and clinical response.

Cattle:

Single-Dose Therapy (BRD Treatment): Administer, by subcutaneous injection, a single dose of 7.5-12.5 mg/kg of body weight (3.4-5.7 mL/100 lb).

Multiple-Day Therapy (BRD Treatment): Administer daily, a subcutaneous dose of 2.5-5 mg/kg of body weight (1.1-2.3 mL/100 lb). Treatment should be repeated at 24-hour intervals for three days. Additional treatments may be given on Days 4 and 5 to animals that have shown clinical improvement but not total recovery.

Single-Dose Therapy (BRD Control): Administer, by subcutaneous injection, a single dose of 7.5 mg/kg of body weight (3.4 mL/100 lb).

Examples of conditions that may contribute to calves being at high risk of developing BRD include, but are not limited to, the following:

- Transportation with animals from two or more farm origins.
- An extended transport time with few to no rest stops.
- An environmental temperature change of $\geq 30^{\circ}\text{F}$ during transportation.
- A $\geq 30^{\circ}\text{F}$ range in temperature fluctuation within a 24-hour period.
- Exposure to wet or cold weather conditions.
- Excessive shrink (more than would be expected with a normal load of cattle).
- Stressful arrival processing procedures (e.g., castration or dehorning).
- Exposure within the prior 72 hours to animals showing clinical signs of BRD.

Administered dose volume should not exceed 20 mL per injection site.

Table 1 – Baytril 100 Dose and Treatment Schedule for Cattle*

| Weight (lb) | Treatment | | Control |
|-------------|---|---|--|
| | Single-Dose Therapy 7.5 - 12.5 mg/kg Dose Volume (mL) | Multiple-Day Therapy 2.5 - 5.0 mg/kg Dose Volume (mL) | Single-Dose Therapy 7.5 mg/kg Dose Volume (mL) |
| 100 | 3.5 - 5.5 | 1.5 - 2.0 | 3.5 |
| 200 | 7.0 - 11.0 | 2.5 - 4.5 | 7.0 |
| 300 | 10.5 - 17.0 | 3.5 - 6.5 | 10.5 |
| 400 | 14.0 - 22.5 | 4.5 - 9.0 | 14.0 |
| 500 | 17.0 - 28.5 | 5.5 - 11.5 | 17.0 |
| 600 | 20.5 - 34.0 | 7.0 - 13.5 | 20.5 |
| 700 | 24.0 - 39.5 | 8.0 - 16.0 | 24.0 |
| 800 | 27.5 - 45.5 | 9.0 - 18.0 | 27.5 |
| 900 | 31.0 - 51.0 | 10.0 - 20.5 | 31.0 |
| 1000 | 34.0 - 57.0 | 11.0 - 23.0 | 34.0 |
| 1100 | 37.5 - 62.5 | 12.5 - 25.0 | 37.5 |

*Dose volumes have been rounded to the nearest 0.5 mL within the dose range.

See product insert for complete dosing and administration information.

Use within 30 days of first puncture and puncture a maximum of 30 times with a needle or 4 times with a dosage delivery device. Any product remaining beyond these parameters should be discarded.

RESIDUE WARNINGS:

Cattle: Animals intended for human consumption must not be slaughtered within 28 days from the last treatment. This product is not approved for female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

HUMAN WARNINGS:

Not for use in humans. Keep out of reach of children. Avoid contact with eyes. In case of contact, immediately flush eyes with copious amounts of water for 15 minutes. In case of dermal contact, wash skin with soap and water. Consult a physician if irritation persists following ocular or dermal exposures. Individuals with a history of hypersensitivity to quinolones should avoid this product. In humans, there is a risk of user photosensitization within a few hours after excessive exposure to quinolones. If excessive accidental exposure occurs, avoid direct sunlight. For product questions, to report adverse reactions, or for a copy of the Safety Data Sheet (SDS), call Elanco Product & Veterinary Support at 1-800-428-4441.

PRECAUTIONS:

The effects of enrofloxacin on cattle reproductive performance, pregnancy and lactation have not been adequately determined.

Subcutaneous injection in cattle can cause a transient local tissue reaction that may result in trim loss of edible tissue at slaughter. Baytril 100 contains different excipients than other Baytril products. The safety and efficacy of this formulation in species other than cattle have not been determined.

Quinolone-class drugs should be used with caution in animals with known or suspected Central Nervous System (CNS) disorders. In such animals, quinolones have, in rare instances, been associated with CNS stimulation which may lead to convulsive seizures. Quinolone-class drugs have been shown to produce erosions of cartilage of weight-bearing joints and other signs of arthropathy in immature animals of various species. See Animal Safety section for additional information.

ADVERSE REACTIONS:

No adverse reactions were observed during clinical trials.

For additional information about reporting adverse drug experiences for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

EFFECTIVENESS:

Cattle: A total of 845 calves with naturally-occurring BRD were treated with Baytril 100 in eight field trials located in five cattle-feeding states. Response to treatment was compared to non-treated controls. Single-dose and multiple-day therapy regimens were evaluated. BRD and mortality were significantly reduced in enrofloxacin-treated calves. No adverse reactions were reported in treated animals.

The effectiveness of Baytril 100 for the control of respiratory disease in cattle at high risk of developing BRD was evaluated in a six-location study in the U.S. and Canada. A total of 1,150 crossbred beef calves at high risk of developing BRD were enrolled in the study. Baytril 100 (7.5 mg/kg BW) or an equivalent volume of sterile saline was administered as a single subcutaneous injection within two days after arrival. Cattle were observed daily for clinical signs of BRD and were evaluated for success on Day 14 post-treatment. Treatment success in the Baytril 100 group (497/573, 87.83%) was significantly higher ($P = 0.0013$) than success in the saline control group (455/571, 80.92%). In addition, there were more treatment successes ($n = 13$) than failures ($n = 3$) in the group of animals positive for *M. bovis* on Day 0 that were treated with Baytril 100. No product-related adverse reactions were reported.

STORAGE CONDITIONS: Protect from direct sunlight. Do not refrigerate or freeze. Store at 20-30°C (68-86°F), excursions permitted up to 40°C (104°F). Precipitation may occur due to cold temperature. To redissolve, warm and then shake the vial.

HOW SUPPLIED:

Baytril 100:
 100 mg/mL 100 mL Bottle
 100 mg/mL 250 mL Bottle
 100 mg/mL 500 mL Bottle

For product questions, to report adverse reactions, or for a copy of the Safety Data Sheet (SDS), call Elanco Product & Veterinary Support at 1-800-428-4441.

Revised: January 2022

Baytril is sold by Elanco or its affiliates and is not a product of Bayer. The product name Baytril is owned by Bayer and used under license. Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates. © 2022 Elanco or its affiliates. PM-US-23-1616

Baytril 100
 Approved by FDA under NADA # 141-068
 Manufactured for: Elanco US Inc.
 Greenfield, IN 46140 U.S.A
 Made in Germany



Loncor™ 300

(florfenicol)



300 mg/mL Injectable Solution

For intramuscular and subcutaneous use in beef and non-lactating dairy cattle only
Not for use in female dairy cattle 20 months of age or older or in calves to be processed for veal

CAUTION

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION

Loncor™ 300 Injectable Solution is a solution of the synthetic antibiotic florfenicol. Each milliliter of sterile Loncor™ 300 contains 300 mg of florfenicol, 250 mg N-Methyl-2-pyrrolidone, 150 mg propylene glycol, and polyethylene glycol qs.

INDICATIONS

Loncor™ 300 is indicated for treatment of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida*, and *Histophilus somni*, and for the treatment of bovine interdigital phlegmon (foot rot, acute interdigital necrobacillosis, infectious pododermatitis) associated with *Fusobacterium necrophorum* and *Bacteroides melaninogenicus*. Also, it is indicated for the control of respiratory disease in cattle at high risk of developing BRD associated with *Mannheimia haemolytica*, *Pasteurella multocida*, and *Histophilus somni*.

DOSAGE AND ADMINISTRATION

For treatment of bovine respiratory disease (BRD) and bovine interdigital phlegmon (foot rot): Loncor™ 300 should be administered by intramuscular injection to cattle at a dose rate of 20 mg/kg body weight (3 mL/100 lbs). A second dose should be administered 48 hours later. Alternatively, Loncor™ 300 can be administered by a single subcutaneous (SC) injection to cattle at a dose rate of 40 mg/kg body weight (6 mL/100 lbs). Do not administer more than 10 mL at each site. The injection should be given only in the neck.

NOTE: Intramuscular injection may result in local tissue reaction which persists beyond 28 days. This may result in trim loss of edible tissue at slaughter. Tissue reaction at injection sites other than the neck is likely to be more severe.

For control of respiratory disease in cattle at high risk of developing BRD: Loncor™ 300 should be administered by a single subcutaneous injection to cattle at a dose rate of 40 mg/kg body weight (6 mL/100 lbs). Do not administer more than 10 mL at each site. The injection should be given only in the neck.

Loncor™ 300 DOSAGE GUIDE

| ANIMAL WEIGHT (lbs) | IM Loncor 300 DOSAGE 3.0 mL/100 lb Body Weight (mL) | SC Loncor 300 DOSAGE 6.0 mL/100 lb Body Weight (mL) |
|---------------------|---|---|
| 100 | 3.0 | 6.0 |
| 200 | 6.0 | 12.0 |
| 300 | 9.0 | 18.0 |
| 400 | 12.0 | 24.0 |
| 500 | 15.0 | 30.0 |
| 600 | 18.0 | 36.0 |
| 700 | 21.0 | 42.0 |
| 800 | 24.0 | 48.0 |
| 900 | 27.0 | 54.0 |
| 1000 | 30.0 | 60.0 |

RECOMMENDED INJECTION LOCATION



Do not inject more than 10 mL per injection site.

Clinical improvement should be evident in most treated subjects within 24 hours of initiation of treatment. If a positive response is not noted within 72 hours of initiation of treatment, the diagnosis should be re-evaluated.

CONTRAINDICATIONS

Do not use in animals that have shown hypersensitivity to florfenicol.

WARNINGS: NOT FOR HUMAN USE. KEEP OUT OF REACH OF CHILDREN.

This product contains materials that can be irritating to skin and eyes. Avoid direct contact with skin, eyes, and clothing. In case of accidental eye exposure, flush with water for 15 minutes. In case of accidental skin exposure, wash with soap and water. Remove contaminated clothing. Consult a physician if irritation persists. Accidental injection of this product may cause local irritation. Consult a physician immediately. The Safety Data Sheet (SDS) contains more detailed occupational safety information. For customer service, adverse effects reporting, and/or a copy of the SDS, call 1-800-422-9874.

PRECAUTIONS

Not for use in animals intended for breeding purposes. The effects of florfenicol on bovine reproductive performance, pregnancy, and lactation have not been determined. Toxicity studies in dogs, rats, and mice have associated the use of florfenicol with testicular degeneration and atrophy. Intramuscular injection may result in local tissue reaction which persists beyond 28 days. This may result in trim loss of edible tissue at slaughter. Tissue reaction at injection sites other than the neck is likely to be more severe.

RESIDUE WARNINGS

Animals intended for human consumption must not be slaughtered within 28 days of the last intramuscular treatment. Animals intended for human consumption must not be slaughtered within 38 days of subcutaneous treatment. This product is not approved for use in female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or calves born to these cows. A withdrawal period has not been established in pre-ruminating calves. Do not use in calves to be processed for veal.

ADVERSE REACTIONS

Inappetence, decreased water consumption, or diarrhea may occur transiently following treatment. To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Elanco at 1-800-422-9874. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

CLINICAL PHARMACOLOGY

The pharmacokinetic disposition of florfenicol was evaluated in feeder calves following single intramuscular (IM) administration at the recommended dose of 20 mg/kg body weight. Florfenicol was also administered intravenously (IV) to the same cattle in order to calculate the volume of distribution, clearance, and percent bioavailability¹ (Table 1).

TABLE 1. Pharmacokinetic Parameter Values for Florfenicol Following IM Administration of 20 mg/kg Body Weight to Feeder Calves (n=10).

| Parameter | Median | Range |
|--------------------------------|--------|-------------|
| C _{max} (µg/mL) | 3.07* | 1.43 - 5.60 |
| T _{max} (hr) | 3.33 | 0.75 - 8.00 |
| T 1/2 (hr) | 18.3** | 8.30 - 44.0 |
| AUC (µg•min/mL) | 4242 | 3200 - 6250 |
| Bioavailability (%) | 78.5 | 59.3 - 106 |
| Vd _{ss} (L/kg)*** | 0.77 | 0.68 - 0.85 |
| Cl _t (mL/min/kg)*** | 3.75 | 3.17 - 4.31 |

* harmonic mean

** mean value

*** following IV administration

C_{max} Maximum serum concentration

T_{max} Time at which C_{max} is observed

T 1/2 Biological half-life

AUC Area under the curve

Vd_{ss} Volume of distribution at steady state

Cl_t Total body clearance

Florfenicol was detectable in the serum of most animals through 60 hours after intramuscular administration with a mean concentration of 0.19 µg/mL. The protein binding of florfenicol was 12.7%, 13.2%, and 18.3% at serum concentrations of 0.5, 3.0, and 16.0 µg/mL, respectively.

MICROBIOLOGY

Florfenicol is a synthetic, broad-spectrum antibiotic active against many Gram-negative and Gram-positive bacteria isolated from domestic animals. It acts by binding to the 50S ribosomal subunit and inhibiting bacterial protein synthesis. Florfenicol is generally considered a bacteriostatic drug, but exhibits bactericidal activity against certain bacterial species. *In vitro* studies demonstrate that florfenicol is active against the bovine respiratory disease (BRD) pathogens *Mannheimia haemolytica*, *Pasteurella multocida*, and *Histophilus somni*, and that florfenicol exhibits bactericidal activity against strains of *M. haemolytica* and *H. somni*. Clinical studies confirm the efficacy of florfenicol against BRD as well as against commonly isolated bacterial pathogens in bovine interdigital phlegmon including *Fusobacterium necrophorum* and *Bacteroides melaninogenicus*.

The minimum inhibitory concentrations (MICs) of florfenicol for BRD organisms were determined using isolates obtained from natural infections from 1990 to 1993. The MICs for interdigital phlegmon organisms were determined using isolates obtained from natural infections from 1973 to 1997 (Table 2).

TABLE 2. Florfenicol Minimum Inhibitory Concentration (MIC) Values* of Indicated Pathogens Isolated From Natural Infections of Cattle.

| Indicated pathogens | Year of isolation | Isolate Numbers | MIC ₅₀ ** (µg/mL) | MIC ₉₀ ** (µg/mL) |
|------------------------------------|-------------------|-----------------|------------------------------|------------------------------|
| <i>Mannheimia haemolytica</i> | 1990 to 1993 | 398 | 0.5 | 1 |
| <i>Pasteurella multocida</i> | 1990 to 1993 | 350 | 0.5 | 0.5 |
| <i>Histophilus somni</i> | 1990 to 1993 | 66 | 0.25 | 0.5 |
| <i>Fusobacterium necrophorum</i> | 1973 to 1997 | 33 | 0.25 | 0.25 |
| <i>Bacteroides melaninogenicus</i> | 1973 to 1997 | 20 | 0.25 | 0.25 |

* The correlation between the *in vitro* susceptibility data and clinical effectiveness is unknown.

** The lowest MIC to encompass 50% and 90% of the most susceptible isolates, respectively.

ANIMAL SAFETY

A 10X safety study was conducted in feeder calves. Two intramuscular injections of 200 mg/kg were administered at a 48-hour interval. The calves were monitored for 14 days after the second dose. Marked anorexia, decreased water consumption, decreased body weight, and increased serum enzymes were observed following dose administration. These effects resolved by the end of the study. A 1X, 3X, and 5X (20, 60, and 100 mg/kg) safety study was conducted in feeder calves for 3X the duration of treatment (6 injections at 48-hour intervals). Slight decrease in feed and water consumption was observed in the 1X dose group. Decreased feed and water consumption, body weight, urine pH, and increased serum enzymes, were observed in the 3X and 5X dose groups. Depression, soft stool consistency, and dehydration were also observed in some animals (most frequently at the 3X and 5X dose levels), primarily near the end of dosing.

A 43-day controlled study was conducted in healthy cattle to evaluate effects of florfenicol administered at the recommended dose on feed consumption. Although a transient decrease in feed consumption was observed, florfenicol administration had no long-term effect on body weight, rate of gain, or feed consumption.

STORAGE INFORMATION

Store below 30°C (86°F).

Once opened, use contents within 6 months.

The solution is light yellow to straw colored. Color does not affect potency.

HOW SUPPLIED

Loncor™ 300 is packaged in 250 mL and 500 mL glass sterile multiple-dose vials.

REFERENCE

1. Lobell RD, Varma KJ, et al. Pharmacokinetics of florfenicol following intravenous and intramuscular doses to cattle. *J Vet Pharmacol Therap.* 1994; 17:253-258.

Made in China

LV2102

© 2021 Elanco or its affiliates.

Approved by FDA under ANADA # 200-582

February, 2021

Loncor™ 300

Loncor, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

Elanco US Inc

Shawnee, KS 66216 U.S.A.



Elanco™

Component™ TE-G

(trenbolone acetate and estradiol implants)

FOR GROWING BEEF STEERS AND HEIFERS ON PASTURE

(Stocker, Feeder, and Slaughter)

20 dose Cartridge Belt

Use with a Component Implanter

DESCRIPTION: Each cartridge belt holds 20 doses of Component TE-G (trenbolone acetate and estradiol implant) Implants. Each dose of 2 pellets consists of 2 pellets containing a total of 40 mg of trenbolone acetate and 8 mg estradiol.

INDICATIONS FOR USE: For increased rate of weight gain in growing beef steers and heifers on pasture (stocker, feeder, and slaughter).

Not approved for repeated implantation (re-implantation) with this or any other cattle ear implant in growing beef steers and heifers on pasture (stocker, feeder, and slaughter). Safety and effectiveness following re-implantation have not been evaluated.

Do not use in beef calves less than 2 months of age, dairy calves, and veal calves because effectiveness and safety have not been established.

Do not use in animals intended for subsequent breeding, or in dairy cows.

Withdrawal Periods and Residue Warnings


No withdrawal period is required when used according to labeling. Do not use in beef calves less than 2 months of age, dairy calves, and veal calves. A withdrawal period has not been established for this product in pre-ruminating calves.

Do not use in dairy cows or in animals intended for subsequent breeding. Use in these cattle may cause drug residues in milk and/or calves born to these cows.

Implant pellets subcutaneously in ear only. Any other location is a violation of Federal law. Do not attempt salvage of implanted site for human or animal food.

Insert the Needle

With one hand firmly grasp the ear. With the other hand insert needle point through the skin and ease forward on a lateral plane until the entire length of the needle is under the skin.



Implant the Pellets

After inserting the needle fully in the correct position, squeeze the trigger fully as the needle is withdrawn from the ear. This properly deposits the implant in the needle track.

This procedure should prevent breakage or crushing of pellets if otherwise forced into contact with tough fibrous-tissue underlying the skin. The length and total contact area of the single dose are designed to permit absorption of the hormones after implantation to stimulate good weight gain. Broken or crushed pellets may interfere with rates of gain.

Storage Conditions

Store unopened product at or below 25°C (77°F). Avoid excessive heat and humidity.

Use before the expiration date printed on foil pouch. Discard open foil pouches.

Approved by FDA under ANADA # 200-221

Manufactured by a non-sterilizing process.

Component, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

© 2023 Elanco or its affiliates

Distributed by Elanco US Inc, Greenfield, IN 46140, USA

QUESTIONS/COMMENTS? For a copy of the Safety Data Sheet or to report side effects, contact Elanco US, Inc. at 1-888-545-5973. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

June 2023

USER SAFETY WARNINGS:

Not for human use. Keep out of reach of children.

Restricted Drug (California) - use only as directed.

IMPLANTING INSTRUCTIONS:

Loading the Implanter

Load the implanter following the instructions supplied with each implanter.

Restrain the Animal

Speed of implantation as well as safety of handlers is best achieved by restraining animal in a squeeze chute using head restraint.




When implanting horned cattle, better control is obtained with additional use of nose tongs.

Prepare the Implant Site

Scrub the back side of the ear (implant site) with a piece of clean absorbent cotton which has been soaked with topical germicidal solution. Follow manufacturer's directions on germicide for correct strength and preparation of solution. Avoid getting disinfectant in animal's eyes.

Where to Implant

The full contents of one cartridge cell should be implanted beneath the skin on the back side of the middle one-third of the ear as illustrated in the drawing. The implant must not be closer to the head than the edge of the auricular cartilage ring farthest from the head. The location for insertion of the needle is a point toward the tip of the ear at least a needle length away from the intended deposition site. Avoid injuring the large arteries, veins and cartilage of the ear.



Elanco™

(progesterone and estradiol benzoate and tylosin tartrate implants)

Each implant consists of 100 mg progesterone USP and 10 mg estradiol benzoate and 29 mg tylosin tartrate

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

IMPLANTS FOR BEEF CALVES 45 DAYS OF AGE AND OLDER AND WEIGHING UP TO 400 LBS**Use with a Component Implanter**

DESCRIPTION: Each cartridge belt holds 20 doses of Component E-C with Tylan (progesterone and estradiol benzoate and tylosin tartrate implant) Implants. Each dose of 5 pellets consists of 4 pellets containing a total of 100 mg progesterone USP and 10 mg estradiol benzoate plus 1 pellet containing 29 mg tylosin tartrate as a local antibacterial.

INDICATIONS FOR USE: For increased rate of weight gain in beef calves 45 days of age and older and weighing up to 400 lbs.

This implant is not approved for repeated implantation (reimplantation) with this or any other cattle ear implant as safety and effectiveness has not been evaluated.

Do not use in calves less than 45 days of age or veal calves because effectiveness and safety have not been evaluated.

Do not use in animals intended for subsequent breeding, or in dairy cows.

DIRECTIONS: Administer 1 implant - the entire contents of one cartridge cell (5 pellets) - subcutaneously in the back of the middle third of the ear. Study and carefully follow at all times the "IMPLANTING INSTRUCTIONS" presented below, avoiding short cuts. Skin infection can be avoided by properly preparing implant site and implanter. During fly season use fly repellent on implant site. One designated team member should always do the implanting. Cleanliness of hands and instruments is important at all times.

Withdrawal Periods and Residue Warnings

No withdrawal period is required when used according to labeling.

Do not use in beef calves less than 45 days of age, dairy calves, and veal calves. Do not use in bull calves intended for reproduction. A withdrawal period has not been established for this product in pre-ruminating calves.

Do not use in dairy cows or in animals intended for subsequent breeding. Use in these cattle may cause drug residues in milk and/or in calves born to these cows. Implant pellets subcutaneously in ear only. Any other location is a violation of Federal law. Do not attempt salvage of implanted site for human or animal food.

USER SAFETY WARNINGS: Not for use in humans. Keep out of reach of children.

Restricted Drug (California) - Use Only as Directed

ANIMAL SAFETY WARNINGS: Bulling has occasionally been reported in implanted steers and heifers. Vaginal and rectal prolapse, udder development, ventral edema and elevated tailheads have occasionally been reported in heifers administered Component E-C with Tylan Implants.

WARNINGS: This product is intended for use in cattle at high risk of developing ear abscesses. Not recommended for use in cattle at low risk of developing ear abscesses (e.g., dry, clean cattle).


The benefit of using a cattle ear implant containing a tylosin tartrate pellet when animals receive a concomitant antimicrobial product has not been evaluated.

IMPLANTING INSTRUCTIONS:**Loading the Implanter**


Load the implanter following the instructions supplied with each implanter.

Restrain the Animal

Speed of implantation as well as safety of handlers is best achieved by restraining animal in a squeeze chute using head restraint.

**Prepare the Implant Site**


Scrub the back side of the ear (implant site) with a piece of clean absorbent cotton or brush which has been soaked with topical germicidal solution. Follow manufacturer's directions on germicide for correct strength and preparation of solution. Avoid getting disinfectant in animal's eyes.

**Where to Implant**

The full contents of one cartridge cell should be implanted beneath the skin on the back side of the middle one-third of the ear as illustrated in the drawing. The implant must not be closer to the head than the edge of the auricular cartilage ring farthest from the head. The location for insertion of the needle is a point toward the tip of the ear and at least a needle length away from the intended deposition site. Avoid injuring the large arteries, veins and cartilage of the ear.

**Insert the Needle**

With one hand firmly grasp the ear. With the other hand insert needle point through the skin and ease forward on a lateral plane until the entire length of the needle is under the skin.

**Implant the Pellets**

After inserting the needle fully in the correct implant position, squeeze the trigger fully as the needle is withdrawn from the ear. This properly deposits the implant in the needle track. This procedure should prevent breakage or crushing of pellets if otherwise forced into contact with tough fibrous-tissue underlying the skin. The length and total contact area of a single dose are designed to permit absorption of the hormones after implantation to stimulate good weight gain. Broken or crushed pellets may lead to undesirable side effects such as bulling, rectal and vaginal prolapse, etc., as noted in the WARNINGS.

STORAGE CONDITIONS: Store at controlled room temperature 15° to 30°C (59° to 86°F) DO NOT refrigerate – avoid excessive heat and humidity. Discard open foil pouches.

Approved by FDA under NADA # 110-315

Manufactured by a non-sterilizing process.

Component E-C with Tylan (progesterone and estradiol benzoate and tylosin tartrate implants) is covered by U.S. Patent No. 5,874,098

Component, Tylan, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

© 2023 Elanco or its affiliates

Distributed by Elanco US Inc., Greenfield, IN 46140, USA

Tylosin: Product of United Kingdom

Revised: June 2023

Elanco™

Component™

TE-IS with Tylan™

(trenbolone acetate and estradiol and tylosin tartrate implants)

FOR GROWING BEEF STEERS FED IN CONFINEMENT FOR SLAUGHTER

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

20-dose Cartridge Belt

Use with a Component Implanter

DESCRIPTION: Each cartridge belt holds 20 doses of Component TE-IS with Tylan (trenbolone acetate and estradiol and tylosin tartrate implant) Implants. Each dose of 5 pellets consists of 4 pellets containing a total of 80 mg of trenbolone acetate and 16 mg estradiol plus 1 pellet containing 29 mg tylosin tartrate as a local antibacterial.

INDICATIONS FOR USE: For increased rate of weight gain and improved feed efficiency in growing beef steers fed in confinement for slaughter.

Not approved for repeated implantation (re-implantation) with this or any other cattle ear implant in growing beef steers fed in confinement for slaughter.

Safety and effectiveness following re-implantation have not been evaluated.

Do not use in beef calves less than 2 months of age, dairy calves, and veal calves because effectiveness and safety have not been established.

Do not use in animals intended for subsequent breeding, or in dairy cows.

WARNING:

This product is intended for use in cattle at high risk of developing ear abscesses. Not recommended for use in cattle at low risk of developing ear abscesses (e.g., dry, clean cattle). The benefit of using a cattle ear implant containing a tylosin tartrate pellet when animals receive a concomitant antimicrobial product has not been evaluated.

Withdrawal Periods and Residue Warnings

No withdrawal period is required when used according to labeling.

Do not use in beef calves less than 2 months of age, dairy calves, and veal calves. A withdrawal period has not been established for this product in pre-ruminating calves.

Do not use in dairy cows or in animals intended for subsequent breeding. Use in these cattle may cause drug residues in milk and/or calves born to these cows.

Implant pellets subcutaneously in ear only. Any other location is a violation of Federal law. Do not attempt salvage of implanted site for human or animal food.

USER SAFETY WARNINGS

Not for human use. Keep out of reach of children.

Restricted Drug (California) - use only as directed.

IMPLANTING INSTRUCTIONS:


Loading the Implanter

Load the implanter following the instructions supplied with each implanter.

Restrain the Animal

Speed of implantation as well as safety of handlers is best achieved by restraining animal in a squeeze chute using head restraint.

When implanting horned cattle, better control is obtained with additional use of nose tongs.



Prepare the Implant Site

Scrub the back side of the ear (implant site) with a piece of clean absorbent cotton which has been soaked with topical germicidal solution. Follow manufacturer's directions on germicide for correct strength and preparation of solution. Avoid getting disinfectant in animal's eyes.

Where to Implant

The full contents of one cartridge cell should be implanted beneath the skin on the back side of the middle one-third of the ear illustrated in the drawing.

The implant must not be closer to the head than the edge of the auricular cartilage ring farthest from the head. The location for insertion of the needle is a point toward the tip of the ear at least a needle length away from the intended deposition site. Avoid injuring the large arteries, veins and cartilage of the ear.

Insert the Needle

With one hand firmly grasp the ear.

With the other hand insert needle point through the skin and ease forward on a lateral plane until the entire length of the needle is under the skin.

Implant the Pellets

After inserting the needle fully in the correct implant position squeeze the trigger fully as the needle is withdrawn from the ear.

This properly deposits the implant in the needle track. This procedure should prevent breakage or crushing of pellets if otherwise forced into contact with tough fibrous-tissue underlying the skin. The length and total contact area of a single dose are designed to permit absorption of the hormones after implantation to stimulate good weight gain. Broken or crushed pellets may interfere with rates of gain.

Storage Conditions

Store unopened product at or below 25°C (77°F). Avoid excessive heat and humidity.

Use before the expiration date printed on the foil pouch. Discard open foil pouches.

Approved by FDA under ANADA # 200-221

Manufactured by a non-sterilizing process.

Component TE-IS with Tylan is covered by U.S. Patent No. 5,874,098

Component, Tylan, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.




© 2023 Elanco or its affiliates

Distributed by Elanco US Inc., Greenfield, IN 46140, USA

Tylosin: Product of the United Kingdom

QUESTIONS/COMMENTS? For a copy of the Safety Data Sheet or to report side effects, contact Elanco US, Inc. at 1-888-545-5973. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

April 2023





(trenbolone acetate and estradiol and tylosin tartrate implants)

FOR GROWING BEEF STEERS FED IN CONFINEMENT FOR SLAUGHTER

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

20-dose Cartridge Belt

Use with a Component Implanter

DESCRIPTION: Each cartridge belt holds 20 doses of Component TE-IS with Tylan (trenbolone acetate and estradiol and tylosin tartrate implant) Implants. Each dose of 5 pellets consists of 4 pellets containing a total of 80 mg of trenbolone acetate and 16 mg estradiol plus 1 pellet containing 29 mg tylosin tartrate as a local antibacterial.

INDICATIONS FOR USE: For increased rate of weight gain and improved feed efficiency in growing beef steers fed in confinement for slaughter.

Not approved for repeated implantation (re-implantation) with this or any other cattle ear implant in growing beef steers fed in confinement for slaughter.

Safety and effectiveness following re-implantation have not been evaluated.

Do not use in beef calves less than 2 months of age, dairy calves, and veal calves because effectiveness and safety have not been established.

Do not use in animals intended for subsequent breeding, or in dairy cows.

WARNING:

This product is intended for use in cattle at high risk of developing ear abscesses. Not recommended for use in cattle at low risk of developing ear abscesses (e.g., dry, clean cattle). The benefit of using a cattle ear implant containing a tylosin tartrate pellet when animals receive a concomitant antimicrobial product has not been evaluated.

Withdrawal Periods and Residue Warnings

No withdrawal period is required when used according to labeling.

Do not use in beef calves less than 2 months of age, dairy calves, and veal calves. A withdrawal period has not been established for this product in pre-ruminating calves.

Do not use in dairy cows or in animals intended for subsequent breeding. Use in these cattle may cause drug residues in milk and/or calves born to these cows.

Implant pellets subcutaneously in ear only. Any other location is a violation of Federal law. Do not attempt salvage of implanted site for human or animal food.

USER SAFETY WARNINGS

Not for human use. Keep out of reach of children.

Restricted Drug (California) - use only as directed.


IMPLANTING INSTRUCTIONS:

Loading the Implanter

Load the implanter following the instructions supplied with each implanter.

Restrain the Animal

Speed of implantation as well as safety of handlers is best achieved by restraining animal in a squeeze chute using head restraint. When implanting horned cattle, better control is obtained with additional use of nose tongs.




Prepare the Implant Site

Scrub the back side of the ear (implant site) with a piece of clean absorbent cotton which has been soaked with topical germicidal solution. Follow manufacturer's directions on germicide for correct strength and preparation of solution. Avoid getting disinfectant in animal's eyes.


Where to Implant

The full contents of one cartridge cell should be implanted beneath the skin on the back side of the middle one-third of the ear illustrated in the drawing. The implant must not be closer to the head than the edge of the auricular cartilage ring farthest from the head. The location for insertion of the needle is a point toward the tip of the ear at least a needle length away from the intended deposition site. Avoid injuring the large arteries, veins and cartilage of the ear.



Insert the Needle

With one hand firmly grasp the ear. With the other hand insert needle point through the skin and ease forward on a lateral plane until the entire length of the needle is under the skin.



Implant the Pellets

After inserting the needle fully in the correct implant position squeeze the trigger fully as the needle is withdrawn from the ear. This properly deposits the implant in the needle track. This procedure should prevent breakage or crushing of pellets if otherwise forced into contact with tough fibrous-tissue underlying the skin. The length and total contact area of a single dose are designed to permit absorption of the hormones after implantation to stimulate good weight gain. Broken or crushed pellets may interfere with rates of gain.

Storage Conditions

Store unopened product at or below 25°C (77°F). Avoid excessive heat and humidity.

Use before the expiration date printed on the foil pouch. Discard open foil pouches.

Approved by FDA under ANADA # 200-221

Manufactured by a non-sterilizing process.

Component TE-IS with Tylan is covered by U.S. Patent No. 5,874,098

Component, Tylan, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

© 2023 Elanco or its affiliates

Distributed by Elanco US Inc., Greenfield, IN 46140, USA

Tylosin: Product of the United Kingdom

QUESTIONS/COMMENTS? For a copy of the Safety Data Sheet or to report side effects, contact Elanco US, Inc. at 1-888-545-5973. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

April 2023

Elanco™

Tylan™ 200 Injection

TM

Professional Size
500 mL

(tylosin injection)

200 mg per mL

For Use In Cattle and Swine Only

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

An Antibiotic

Use automatic syringe equipment only

Indications: In Beef Cattle and Non-lactating Dairy Cattle, Tylan 200 Injection is indicated for use in the treatment of bovine respiratory complex (shipping fever, pneumonia) usually associated with *Pasteurella multocida* and *Arcanobacterium pyogenes*; foot rot (necrotic pododermatitis) and calf diphtheria caused by *Fusobacterium necrophorum* and metritis caused by *Arcanobacterium pyogenes*.

In Swine, Tylan 200 Injection is indicated for use in the treatment of swine arthritis caused by *Mycoplasma hyosynoviae*; swine pneumonia caused by *Pasteurella* spp.; swine erysipelas caused by *Erysipelothrix rhusiopathiae*; swine dysentery associated with *Treponema hyodysenteriae* when followed by appropriate medication in the drinking water and/or feed.

Each mL contains 200 mg of tylosin activity (as tylosin base) in 50 percent propylene glycol with 4 percent benzyl alcohol and water for injection.

ADMINISTRATION AND DOSAGE: Tylan 200 Injection is administered intramuscularly.

BEEF CATTLE AND NON-LACTATING DAIRY CATTLE—Inject intramuscularly 8 mg per pound of body weight one time daily (1 mL per 25 pounds). Treatment should be continued 24 hours following remission of disease signs, not to exceed 5 days. Do not inject more than 10 mL per site.

SWINE—Inject intramuscularly 4 mg per pound of body weight (1 mL per 50 pounds) twice daily. Treatment should be continued 24 hours following remission of disease signs, not to exceed 3 days. Do not inject more than 5 mL per site.

Read accompanying directions fully before use.

CAUTION:

Do not mix Tylan 200 Injection with other injectable solutions as this may cause a precipitation of the active ingredients.

WARNINGS:

NOT FOR HUMAN USE. KEEP OUT OF REACH OF CHILDREN.

Adverse reactions, including shock and death may result from overdose in baby pigs.

Do not attempt injection into pigs weighing less than 25 pounds (0.5 mL) with the common syringe. It is recommended that Tylan 50 Injection be used in pigs weighing less than 25 pounds.

Do not administer to horses or other equines. Injection of tylosin in equines has been fatal.

RESIDUE WARNING: Swine: Swine intended for human consumption must not be slaughtered within 14 days of the last use of this drug product.

RESIDUE WARNING: Cattle: Cattle intended for human consumption must not be slaughtered within 21 days of the last use of this drug product. This drug product is not approved for use in female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows. This product is not approved for use in calves intended to be processed for veal. A withdrawal period has not been established in pre-ruminating calves.

If tylosin medicated drinking water is used as a follow-up treatment for swine dysentery, the animal should thereafter receive feed containing 40 to 100 grams of tylosin per ton for 2 weeks to assure depletion of tissue residues.

Store at or below 25°C (77°F).

Tylan, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

© 2023 Elanco or its affiliates. PM-US-23-1624

Approved by FDA under NADA # 012-965

For additional information about reporting adverse drug experiences for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

Manufactured for: Elanco US Inc.

Greenfield, IN 46140, USA

Product of Ireland