

25 Doses
25 Vials of Vaccine
plus 25 x 0.5 mL Vials of Diluent

Canine Distemper- Adenovirus Type 2- Coronavirus-Parvovirus Vaccine Modified Live and Killed Virus Leptospira Canicola-Grippytyphosa- Icterohaemorrhagiae-Pomona Bacterial Extract TruCan™ Ultra DAP+CL4

For use in animals only

This product has been shown to be effective for the vaccination of healthy dogs 6 weeks of age or older against canine distemper, infectious canine hepatitis, canine adenovirus type 2, canine coronavirus, canine parvovirus (CPV), *Leptospira canicola*, *Leptospira grippytyphosa*, *Leptospira icterohaemorrhagiae* and *Leptospira pomona*. The duration of immunity for this product has not been determined. For more information regarding efficacy and safety data, go to productdata.aphis.usda.gov.

This product contains a CPV 2b strain which has been demonstrated effective against disease caused by CPV 2c in puppies with CPV maternal antibody.

Directions for use

- **General directions:** Aseptically rehydrate TruCan Ultra DAP with low volume diluent supplied. Administer one 0.5 mL dose subcutaneously.
- **Primary vaccination:** The presence of maternal antibody is known to interfere with the development of active immunity in dogs and additional boosters will be required in most young animals. All dogs should initially receive one dose of this product and a second dose 2 to 3 weeks later.

Historically, annual revaccination with a single dose has been recommended for this product. The need for this booster has not been established. For advice on revaccination frequency and annual booster vaccinations, consultation with a veterinarian is recommended.

Caution: In the absence of a veterinarian-client-patient relationship, Federal law prohibits the relabeling, repackaging, resale, or redistribution of the individual contents of this package.

Caution: Store in the dark at 2° to 8°C (35° to 46°F). AVOID FREEZING. SHAKE WELL. Do not mix with other products, except as specified on this label. This product has not been tested in pregnant animals. In case of anaphylactoid reaction, administer epinephrine. Inactivate all unused contents before disposal. In case of human exposure, contact a physician.

Gentamicin and thimerosal added as preservatives.

Elanco US Inc.

Fort Dodge, IA 50501

Phone: 888-545-5973

VLN/PCN 196/45J9.20

TruCan, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

© 2021 Elanco or its affiliates.

