



226

BTO2-REO

**DESCRIPTION:**

Bursal disease (standard and variant), Reovirus vaccine, killed virus

DIRECTIONS FOR USE:

Vaccine should be warmed to 32° – 35°C (90° – 95°F) before use. Shake well before using. Inject subcutaneously using aseptic technique. Use entire contents of bottle when first opened. Vaccinate (prime) with live virus tenosynovitis and bursal disease vaccines at least 4 weeks prior to the use of this vaccine. Vaccinate between 12 and 20 weeks of age. Revaccinate during molt.

DOSE:

1/2 ml (0.5 ml) per bird. Inject all birds in flock.

RECOMMENDED USE:

Recommended for the vaccination of breeder hens and replacement pullets as an aid to provide maternal antibodies for the early protection of progeny against Reovirus related malabsorption syndrome and tenosynovitis and to type 1, infectious bursal disease, standard and variant strains. Aids in the reduction of BD AL-2 Variant in chickens. BD AL-2 Variant protection has not been demonstrated in progeny.

CAUTION:

Store at 2° - 8°C (35° - 46°F). Do not freeze.

WARNING:

In case of accidental human injection, seek immediate medical attention. Do not vaccinate within 42 days before slaughter. Contact Elanco with questions or concerns at 1-800-428-4441 or 1-888-545-5973.

LIST NO: KV: 7800

FOR VETERINARY USE ONLY

Further information: www.elanco.us



The label contains complete use information, including cautions and warnings. Always read, understand and follow the label and use directions.

AviPro, Elanco, and the diagonal bar logo are trademarks of Elanco or its affiliates. © 2019 Elanco. PM-US-19-1564(2)



1-800-428-4441
www.elanco.us