

AviPro®

329 ND-IB2-SE4 Conc

Description: Newcastle disease, bronchitis vaccine (Mass. and Ark. type), killed virus, *Salmonella* Enteritidis bacterin

Directions For Use: Vaccine should be warmed to room temperature before use. Shake well before using. Inject subcutaneously or intramuscularly using aseptic technique. Use entire contents of bottle when first opened. Vaccinate (prime) with live virus vaccines for Newcastle disease and bronchitis at least 4 weeks prior to the use of this vaccine. Vaccinate subcutaneously at 12 and 16 weeks of age or intramuscularly at 13 and 17 weeks of age. Revaccinate with monovalent *Salmonella* Enteritidis bacterin 4 weeks later. Revaccinate during molt.

Dose: 1/4 ml (0.25 ml) per bird.
Inject all birds in flock.

The label contains complete use information, including cautions and warnings. Always read, understand and follow the label and use directions.

Elanco®, AviPro®, Full Value Poultry™ and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries, or affiliates.

© 2016 Eli Lilly and Company, its subsidiaries or affiliates.
USPBVAVP00032

Recommended for the vaccination of chickens as an aid in the prevention of Newcastle disease and Infectious Bronchitis (Mass. and Ark. types) and as an aid to reduce the colonization of the internal organs, including the reproductive tract, by *Salmonella* Enteritidis.

Caution: Store at 2°-7°C (35°-45°F). Do not freeze.

Warning: In case of accidental human injection seek immediate medical attention. Do not vaccinate within 42 days before slaughter. Contact Elanco with questions or concerns at 1-800-428-4441 or 1-888-545-5973.

LISTNO: KV: 8811-H

FOR VETERINARY USE ONLY
Further information: www.elanco.us




FULL VALUE POULTRY™

Elanco™

1-800-428-4441
www.elanco.us