

AVIPRO[®]

109 SE4 Conc

VACCINES



Description

Salmonella Enteritidis Bacterin

Directions for Use:

Bacterin should be warmed to room temperature before use. Shake well before using. Inject subcutaneously using aseptic technique. Use entire contents of bottle when first opened. Vaccinate chickens between 12 and 16 weeks of age. Revaccinate four weeks later. Revaccinate during molt.

Dose:

1/4 ml (0.25 ml) per bird. Inject all birds in flocks.

Follow directions carefully. As with any oil emulsion bacterin, improper vaccination technique may result in a reaction at the site of injection. Do not administer intramuscularly.

Strain Information:

Salmonella Enteritidis phage types 8, 14b, 23 and 24



Recommended for the vaccination of chickens as an aid in the reduction of *Salmonella* Enteritidis colonization of internal organs, including the reproductive tract.

Caution:

Store at 2°–8°C (35°–46°F). Do not freeze.

Warning:

In case of accidental human injection, seek immediate medical attention. Do not vaccinate within 42 days before slaughter. Contact Elanco with questions or concerns at 1-800-428-4441 or 1-888-545-5973.

LIST NO:

KV: 8809-H

FOR VETERINARY USE ONLY

For more information, visit elanco.us/poultry

The label contains complete use information, including cautions and warnings. Always read, understand and follow the label and use directions.

AviPro, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.
© 2020 Elanco. PM-US-20-0479

AviPro[®]



Poultry