



IB ARK



DESCRIPTION:

Bronchitis vaccine (Ark. type), live virus

DIRECTIONS FOR USE:

Vaccinate chickens at 1 day of age or older using the drinking water, intraocular or coarse spray method. A second dose should be given at 2 weeks of age or older. Replacement pullets should be revaccinated between 16 and 18 weeks of age.

DRINKING WATER METHOD:

Remove any disinfectant or sanitizers from the drinking water for 24 hours before and after vaccination. Rinse all waterers with clean water. Do not use disinfectants. Withhold all water for at least one hour prior to giving vaccine. Mix vaccine in the drinking water according to the following schedule:

Liters of water to be used:

Age of Birds	1,000 Doses	5,000 Doses	10,000 Doses	20,000 Doses
Under 3 weeks	10	50	100	200
3 to 8 weeks	20	100	200	400
Over 8 weeks	40	200	400	800

Provide ample water space so that all birds can drink easily. Administration through the water lines with a proportioner or medication tank is not recommended.

INTRAOCULAR (EYE DROP) METHOD:

Specific for 100 and 1,000 dose sizes. Rehydrate the vaccine with the diluent provided. Hold the chick with one eye turned up. Put one drop in eye and hold until bird swallows.

COARSE SPRAY METHOD:

- Specific for 2,000, 2,500, 3,000, 5,000, 10,000, and 20,000 dose sizes.
- Rehydrate with cool distilled water in an amount sufficient to vaccinate the required number of chickens according to the directions of the spray cabinet or field sprayer manufacturer

- Field spray application should be used only in houses that can be tightly closed
- Close all windows, doors, ventilators, etc., and turn off fans
- Spray each 1,000 doses of vaccine directly into the faces of each 1,000 birds
- Leave the house closed for at least 15 minutes after spraying
- Always use eye goggles and a respirator

Recommended for the vaccination and revaccination of chickens as an aid in the prevention of infectious bronchitis, Arkansas type.

CAUTION:

Store at 2°-7°C (35° - 45°F). Do not freeze.

WARNING:

Do not vaccinate within 21 days before slaughter. Contains penicillin and streptomycin or gentamicin as preservatives. Contact Elanco with questions or concerns at 1-800-428-4441 or 1-888-545-5973.

LIST NO: LV:6818-K

FOR VETERINARY USE ONLY

Further information: www.elanco.us


The label contains complete use information, including cautions and warnings.

Always read, understand and follow the label and use directions.

AviPro, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

© 2019 Elanco or its affiliates.

PM-US-19-0366



1-800-428-4441
www.elanco.us