



ELSEVIER

Contents lists available at ScienceDirect

## Journal of Affective Disorders

journal homepage: [www.elsevier.com/locate/jad](http://www.elsevier.com/locate/jad)

## Research paper

Association of *CRTC1* polymorphisms with obesity markers in subjects from the general population with lifetime depression

Lina Quteineh<sup>a</sup>, Martin Preisig<sup>b</sup>, Margarita Rivera<sup>c,d,e</sup>, Yuri Milaneschi<sup>f</sup>, Enrique Castelao<sup>b</sup>, Mehdi Gholam-Rezaee<sup>b</sup>, Frederik Vandenberghe<sup>a</sup>, Nuria Saigi-Morgui<sup>a</sup>, Aurélie Delacrétaz<sup>a</sup>, Jean-René Cardinaux<sup>g,h</sup>, Gonneke Willemssen<sup>i</sup>, Dorret I. Boomsma<sup>i</sup>, Brenda W.J.H. Penninx<sup>f</sup>, Ana Ching-López<sup>d</sup>, Philippe Conus<sup>j</sup>, Chin B. Eap<sup>a,k,\*</sup>

<sup>a</sup> Unit of Pharmacogenetics and Clinical Psychopharmacology, Centre for Psychiatric Neuroscience, Department of Psychiatry, Lausanne University Hospital, Prilly, Switzerland

<sup>b</sup> Centre of Psychiatric Epidemiology and Psychopathology, Department of Psychiatry, Lausanne University Hospital, Prilly, Switzerland

<sup>c</sup> Instituto de Investigación Biosanitaria ibs.GRANADA, Hospitales Universitarios de Granada/Universidad de Granada, Granada, Spain

<sup>d</sup> CIBERSAM–University of Granada and Institute of Neurosciences Federico Olóriz, Centro de Investigación Biomédica, University of Granada, Spain

<sup>e</sup> MRC Social Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, United Kingdom

<sup>f</sup> Department of Psychiatry, EMGO Institute for Health and Care Research and Neuroscience Campus Amsterdam, VU University Medical Center/GGZ inGeest, Amsterdam, The Netherlands

<sup>g</sup> Centre for Psychiatric Neuroscience, Department of Psychiatry, Lausanne University Hospital, Prilly, Switzerland

<sup>h</sup> Child and Adolescent Psychiatric Clinic, Department of Psychiatry, Lausanne University Hospital, Prilly, Switzerland

<sup>i</sup> Department of Biological Psychology, VU University Amsterdam, Amsterdam, The Netherlands

<sup>j</sup> Service of General Psychiatry, Department of Psychiatry, Lausanne University Hospital, Prilly, Switzerland

<sup>k</sup> School of Pharmaceutical Sciences, University of Geneva, University of Lausanne, Geneva, Switzerland

## ARTICLE INFO

## Article history:

Received 23 November 2015

Received in revised form

10 February 2016

Accepted 9 March 2016

Available online 10 March 2016

## Keywords:

Obesity

Major depressive disorder

Psychiatric disorders

Psychotropic drugs

Pharmacogenetics

Genetic polymorphisms

## ABSTRACT

**Background:** Psychiatric disorders have been hypothesized to share common etiological pathways with obesity, suggesting related neurobiological bases. We aimed to examine whether *CRTC1* polymorphisms were associated with major depressive disorder (MDD) and to test the association of these polymorphisms with obesity markers in several large case-control samples with MDD.

**Methods:** The association between *CRTC1* polymorphisms and MDD was investigated in three case-control samples with MDD (PsyCoLaus  $n_1=3,362$ , Radiant  $n_2=3,148$  and NESDA/NTR  $n_3=4,663$ ). The effect of *CRTC1* polymorphisms on obesity markers was then explored.

**Results:** *CRTC1* polymorphisms were not associated with MDD in the three samples. *CRTC1* rs6510997C > T was significantly associated with fat mass in the PsyCoLaus study. In fact, a protective effect of this polymorphism was found in MDD cases ( $n=1,434$ ,  $\beta=-1.32\%$ , 95% CI  $-2.07$  to  $-0.57$ ,  $p<0.001$ ), but not in controls. In the Radiant study, *CRTC1* polymorphisms were associated with BMI, exclusively in individuals with MDD ( $n=2,138$ ,  $\beta=-0.75$  kg/m<sup>2</sup>, 95% CI  $-1.30$  to  $-0.21$ ,  $p=0.007$ ), while no association with BMI was found in the NESDA/NTR study.

**Limitations:** Estimated fat mass using bioimpedance that capture more accurately adiposity was only present in the PsyCoLaus sample.

**Conclusions:** *CRTC1* polymorphisms seem to play a role with obesity markers in individuals with MDD rather than non-depressive individuals. Therefore, the weak association previously reported in the population-based samples was driven by cases diagnosed with lifetime MDD. However, *CRTC1* seems not to be implicated directly in the development of psychiatric diseases.

© 2016 Elsevier B.V. All rights reserved.

## 1. Introduction

Based on a growing body of evidence, it has been hypothesized that psychiatric disorders, such as schizophrenia and mood disorders, are causally related to or share common etiological pathways with obesity, suggesting that comorbid obesity and psychiatric disorders

\* Correspondence to: Hôpital de Cery, 1008 Prilly, Lausanne, Switzerland.

E-mail address: [chin.eap@chuv.ch](mailto:chin.eap@chuv.ch) (C.B. Eap).

have related neurobiological bases (Farmer et al., 2008; Forty et al., 2014). Increased appetite, weight gain and obesity are common symptoms of depression (Blaine, 2008; Luppino et al., 2010). On the other hand, obese individuals were also found to have higher prevalence of depression (Luppino et al., 2010). Shared genetic factors were previously suggested (Afari et al., 2010; Comings et al., 1996; Rivera et al., 2012; Samaan et al., 2013), however, due to the pathological complexity of both conditions and the polygenic factors affecting them, many genetic factors are still to unravel. Additionally, due to heterogeneity of depression, genetic association studies could give conflicting results. One example is the fat mass and obesity-associated gene (*FTO*) for which the common *rs9939609-A* variant previously associated with increased body mass index (BMI) and obesity in several studies (Frayling et al., 2007; Qi et al., 2008; Scuteri et al., 2007) was shown to be negatively associated with depression in a meta-analysis of four large studies (Samaan et al., 2013). On the other hand, the same variant was found to be positively associated with depression in a recent large case-control study (Milaneschi et al., 2014), but this association was entirely driven by the atypical depression subtype (subtype characterized typically by increased appetite, weight gain and hypersomnia).

We previously showed an association between a coding single nucleotide polymorphism (SNP) within the *CREB-regulated transcription coactivator 1 (CRTC1)* gene (*rs3746266A > G*) and BMI in 3 independent psychiatric samples (brief description of these samples is given in the Supplementary data and is described in details in ref (Choong et al., 2013)). Carriers of the *rs3746266-G* variant showed significantly protective effect against obesity compared to non-carriers. The sex-stratified analysis in the 3 combined psychiatric samples showed a protective effect for the *G* allele both in men and women. However, the strongest and most clinically relevant association was observed in women younger than 45 years old. Although the effect was weaker, we also found a protective effect of the *T* allele of the *CRTC1 rs6510997C > T* (a proxy of *rs3746266A > G*,  $r^2=0.7$ ) on fat accumulation in a large population-based sample (CoLaus), with the strongest association again in premenopausal women (Choong et al., 2013).

In animal models, we previously showed that mice lacking *CRTC1* exhibit neurobehavioral endophenotypes related to mood disorders, depression-related behavior and a blunted behavioral response to antidepressants (Breuillaud et al., 2012). *Crtc1* knockout mice also developed obese features, including obesity-related metabolic complications, under normal diets (Altarejos et al., 2008; Breuillaud et al., 2009). Altogether, these findings suggest that the *Crtc1* gene might play a common role in obesity and depression-related behavior.

In the present study, we aimed to examine whether *CRTC1* SNPs are associated with major depressive disorder (MDD), and to test the association of *CRTC1* SNPs with obesity markers in several large case-control samples with MDD. The same *CRTC1* SNPs (*rs3746266A > G* and *rs6510997C > T*) investigated previously (Choong et al., 2013) were used in the current study.

## 2. Patients and methods

The associations between *CRTC1* polymorphisms and obesity markers were first tested in the PsyCoLaus sample and tested for replication in the Radiant and NESDA/NTR study. Supplementary Table S1 presents the main similarities and differences between these samples.

## 3. Discovery cohort: CoLaus|PsyCoLaus

Participants aged 35–75 years in this population-based cohort study (CoLaus;  $n=6,734$ ) were recruited between June 2003 and

May 2006 in Lausanne, Switzerland as previously described (Firmann et al., 2008). The assessments at baseline and at the first follow-up included cardiovascular risk factors such as BMI, fat mass, blood pressure, glucose and lipid profiles. In addition, a genome-wide association study was performed in all Caucasians (91% of the sample). All participants of CoLaus were asked to also participate in the psychiatric evaluations (PsyCoLaus (Preisig et al., 2009)), which included the semi-structured Diagnostic Interview for Genetic Studies (DIGS) (Nurnberger et al., 1994; Preisig et al., 1999). The assignment of the lifetime diagnosis of MDD and the subtyping into atypical and non-atypical MDD was performed according to DSM-IV criteria. Complete physical and psychiatric data were available for 3354 participants aged 35–65 year-old who completed the psychiatric evaluation either at the baseline or the first follow-up ( $n_{\text{total}}=3362$ ). 1434 cases of lifetime depression were compared with 1920 controls with no previous history of depression. Genotyping for the CoLaus/PsyCoLaus subjects was performed using the Affymetrix GeneChipR Human Mapping 500 K array set. The *CRTC1 rs6510997C > T* SNP was directly genotyped (Choong et al., 2013). The *CRTC1 rs7257846T > C*, which is in linkage disequilibrium (LD) with *rs3746266A > G* ( $r^2=0.93$ ), was imputed and analyzed ( $r^2$ hat value was 0.70).

The Institutional Ethics Committee of the University of Lausanne approved the CoLaus and subsequently the PsyCoLaus study. All participants signed a written informed consent to participate in the study.

### 3.1. The radiant study

The Radiant study was founded from three studies: the Depression Case-Control (DeCC) study, Depression Network (DeNT) study and the Genome-Based Therapeutic Drugs for Depression (GENDEP) study. The DeCC study is a multicentre case-control study conducted over three investigative sites in UK (London, Cardiff and Birmingham) (Cohen-Woods et al., 2009). The DeNT study is a family-based study that recruited sibling pairs affected with unipolar depression from eight European clinical sites and one in the USA (Farmer et al., 2004; McGuffin et al., 2005). Participants in the DeCC and DeNT studies were included if they had lifetime diagnosis of two or more episodes of MDD of at least moderate severity. Subjects in the GENDEP study were recruited from nine European centers if they experienced at least one depressive episode of at least moderate severity (Uher et al., 2009). Diagnosis of MDD was defined as of illness fulfilling ICD-10 and/or DSM-IV criteria and was ascertained using the Schedules for Clinical Assessment in Neuropsychiatry (SCAN) interview in all three studies (Wing et al., 1990). Control subjects were recruited via the Medical Research Council general practice research framework (Sham et al., 2000). They were screened for lifetime absence of any psychiatric disorder using a modified version of the Past History Schedule (McGuffin et al., 1986). Self-reported height and weight were obtained from participants in the MDD group via the SCAN interview and from the controls via telephone interview. The reliability of self-report of height and weight was assessed in the GENDEP dataset ( $n=811$ ) where also height and weight were measured. The correlations for measured versus self-reported height, weight and BMI were 0.97, 0.95 and 0.95, respectively (Hung et al., 2014, 2015). All cases and controls were of white European ancestry. Genotyping for the Radiant subjects was performed using the Illumina HumanHap610-Quad BeadChips by the Centre National de Génotypage (CNG), Evry, France, as previously described (Lewis et al., 2010). *CRTC1 rs3746266A > G* was successfully imputed using the Beagle Program (BEAGLE Version 3.0 Copyright (c) 2007–2010 Brian L Browning), with imputation quality  $> 0.8$ . The *CRTC1 rs2075017T > C*, which is in complete linkage disequilibrium (LD) with the *rs6510997C > T*, was also

selected for the association analyses.

Approval was obtained from the local research ethics committees/institutional research boards of all of the participating sites. All participants provided written informed consent.

### 3.2. NESDA/NTR study

This sample consisted of 4663 unrelated participants of European ancestry from the Netherlands Study of Depression and Anxiety (NESDA) (Penninx et al., 2008) (1636 cases and 425 controls) and from the Netherlands Twin Registry (NTR) (Boomsma et al., 2006) (132 cases and 2470 controls). The diagnosis of lifetime and/or current MDD according to DSM-IV was ascertained using the Composite Interview Diagnostic Instrument (WHO WMH-CIDI, 2015). Weight and height were measured at the study clinic during the visit for NESDA (Penninx et al., 2008), and during the home visit for NTR-Biobank (Willemsen et al., 2010). Genotyping was performed on multiple chip platforms in (partially overlapping) different subsets of the total sample (Affymetrix-Perlegen 5.0, Illumina 370 K, Illumina 660 K, Illumina Omni 1 M and Affymetrix 6.0) and data were imputed using the 1000 Genomes phase 1 INTEGRATED RELEASE version 3 ALL panel ( $r^2$  hat value was 0.99 for *rs6510997C > T* and 0.77 for *rs3746266A > G*).

The NESDA/NTR study was approved by the Central Ethics Committee on Research Involving Human Subjects of the VU University Medical Center, Amsterdam, and all subjects provided written informed consent.

## 4. Statistical analysis

For association studies, chi-square ( $\chi^2$ ) or Fisher exact tests for binomial variables were used. Differences in genotype frequencies as well as deviation from Hardy-Weinberg equilibrium were assessed using  $\chi^2$  test. All genetic analyses for *CRTC1* SNPs were performed using a dominant model (wild type vs. variant allele carriers), the same model used for the previously mentioned psychiatric samples (Choong et al., 2013).

Multivariable regression analyses were used to test the association between the *CRTC1* SNPs and MDD adjusting for age, sex and BMI. For PsyCoLaus data, a Generalized Linear Model (GLM) adjusted for age and sex was used to test the association between *CRTC1 rs6510997C > T* polymorphism and obesity markers (BMI, fat mass and waist circumference) using SAS 9.3 version (SAS Institute Inc., Cary, NC, USA). In a first step, we tested for the interaction between *CRTC1* SNPs and MDD status for an effect on obesity markers. For the Radiant and NESDA/NTR studies BMI was the only available obesity marker. For the data of these studies, linear regression models adjusted for age, sex and principal components were used to test the association between *CRTC1* polymorphisms and BMI.

## 5. Results

### 5.1. Demographic characteristics of the study samples

General characteristics of the PsyCoLaus sample are presented in Supplementary Table S2. The Radiant study included 3148 subjects (2338 cases and 810 controls), mean age  $\pm$  standard deviation (s.d.) was  $43.9 \pm 12.8$  years. Females represent nearly 68% of the study sample. Mean BMI in the total study sample was  $25.9 \text{ kg/m}^2$  (s.d. 5.4). The NESDA/NTR study included 4663 subjects, mean age  $\pm$  s.d. was  $42.7 \pm 14.0$  years. Females represent nearly 64% of the study sample. Mean BMI in the total study sample was  $25.0 \text{ kg/m}^2$  (s.d. 4.5).

**Table 1**

Genotype distribution in the total study samples, among MDD subjects and controls.

Patients	Total	N (%)			HWE
<b>PsyCoLaus</b>					
<b><i>rs6510997C &gt; T</i><sup>a</sup></b>					
Total	3347	CC	CT	TT	0.37
MDD cases	1431	912 (63.7)	468 (32.7)	51 (3.6)	
Controls	1916	1216 (63.5)	624 (32.6)	76 (3.9)	
<b>The radiant study</b>					
<b><i>rs3746266A &gt; G</i></b>					
Total	2842	AA	AG	GG	0.78
MDD cases	2142	1637 (76.4)	468 (21.9)	37 (1.7)	
Controls	700	536 (76.5)	158 (22.6)	6 (0.9)	
<b><i>rs2075017T &gt; C</i><sup>a</sup></b>					
Total	2160	TT	TC	CC	0.30
MDD cases	1352	849 (62.8)	450 (33.3)	53 (3.9)	
Controls	808	508 (62.9)	270 (33.4)	30 (3.7)	
<b>NESDA/NTR study</b>					
<b><i>rs3746266A &gt; G</i></b>					
Total	4663	AA	AG	GG	0.47
MDD cases	1768	1234 (69.8)	485 (27.4)	49 (2.8)	
Controls	2895	2001 (69.1)	821 (28.5)	73 (2.5)	
<b><i>rs6510997C &gt; T</i></b>					
Total	4663	CC	CT	TT	0.68
MDD cases	1768	1124 (63.6)	567 (32.1)	77 (4.4)	
Controls	2895	1828 (63.1)	955 (33.0)	112 (3.9)	

MDD: Major depressive disorder, N: number, HWE: Hardy-Weinberg Equilibrium.

<sup>a</sup> Both SNPs are in complete linkage disequilibrium ( $r^2=1$ ).

### 5.2. *CRTC1* polymorphisms and MDD

Genotype distribution in the overall study samples and among MDD cases and controls are presented in Table 1. Genotype distribution did not differ between MDD cases and controls in any of the three studies (Table 1). Multivariate analyses adjusted for age and sex confirmed the absence of an association between *CRTC1* polymorphisms and MDD in the three studies (data not shown).

### 5.3. *CRTC1* polymorphisms and obesity markers

#### 5.3.1. PsyCoLaus

*CRTC1 rs6510997C > T* was significantly associated with fat mass in the global PsyCoLaus sample (Table 2). We observed interactions between *CRTC1 rs6510997C > T* and sex ( $p=0.03$ ) as well as between this SNP and MDD status ( $p=0.01$ ) regarding the fat mass, indicating a significantly stronger effect of this genetic variant in females and in subjects with MDD. Indeed, this SNP was significantly associated with lower fat mass in females but not in males. Similarly, this SNP was significantly associated with lower fat mass in subjects with MDD but not in controls (Table 2). Additionally, the 3-way interaction between *CRTC1 rs6510997C > T*, sex and MDD regarding the fat mass was not significant ( $p=0.54$ ). In females, we did not find an interaction between the *CRTC1 rs6510997C > T* SNP and menopausal status and among depressed subjects there was no interaction between this SNP and the atypical depression subtype to affect the fat mass. Regarding the BMI and waist circumference there was no evidence of an interaction between *CRTC1 rs6510997C > T* and sex or MDD status and the SNP was not associated with these obesity markers (Table 2).

No difference on the associations between *CRTC1 rs6510997C > T* and fat mass was noticed by adjusting the GLM model for other available co-variables that could affect the gene association with obesity markers, like socioeconomic status, drug dependence, alcohol

**Table 2**  
Association between *CRTC1* rs6510997C > T polymorphism and fat mass, BMI and waist circumference in the PsyCoLaus sample, among sex-stratified and MDD subjects and controls.

	n	Fat mass <sup>a,b</sup>		BMI <sup>c</sup>		Waist circumference <sup>c</sup>	
		Estimates (95% C.I.)	p-Value	Estimates (95% C.I.)	p-Value	Estimates (95% C.I.)	p-Value
<b>All subjects</b>	3362	−0.66 (−1.28 to −0.04)	<b>0.04</b>	−0.28 (−0.60 to 0.04)	0.09	−0.70 (−1.64 to 0.24)	0.14
Males	1576	−0.03 (−0.56 to 0.51)	0.92				
Females	1786	−1.08 (−1.81 to −0.34)	<b>0.004</b>				
<b>Subjects with MDD</b>	1434	−1.32 (−2.07 to −0.57)	<b>&lt; 0.001</b>				
<b>Controls</b>	1920	−0.01 (−0.60 to 0.57)	0.97				

Models were adjusted for age and sex (when appropriate).

BMI: body mass index, C.I.: confidence interval, MDD: Major depressive disorder.

<sup>a</sup> *CRTC1* rs6510997C > T and MDD status interaction regarding fat mass: p=0.014.

<sup>b</sup> *CRTC1* rs6510997C > T and sex interaction regarding fat mass: estimate: p=0.026.

<sup>c</sup> *CRTC1* rs6510997C > T and MDD status and/or sex interactions regarding BMI and waist circumference: p > 0.05.

consumption, former and current tobacco consumer (Supplementary Table S3). Nearly the same associations with obesity markers were observed in PsyCoLaus for *CRTC1* rs7257846T > C (in strong LD with rs3746266A > G,  $r^2 = 0.93$ ) (Supplementary Table S4).

### 5.3.2. The radiant and NESDA/NTR studies

The *CRTC1* rs3746266A > G SNP was associated with BMI in the total Radiant sample and exclusively in females (Table 3). By stratifying MDD cases and controls, the protective effect of the rs3746266G-allele was only observed in MDD cases, while no association was observed in controls. A weak association was observed between *CRTC1* rs2075017T > C (in complete LD with the rs6510997C > T) and BMI in the Radiant sample (Table 3). Regarding the NESDA/NTR study, *CRTC1* SNPs were not associated with BMI (Table 4).

## 6. Discussion

We previously showed a protective effect of the variant *CRTC1* rs3746266G-allele on increased BMI in three independent psychiatric samples (Choong et al., 2013). The same protective effect was also observed in a population-based sample (CoLaus) between the rs6510997T-allele (in a strong LD with the rs3746266G-allele)

**Table 3**  
Association between *CRTC1* polymorphisms and BMI in the total, sex-stratified sample and in depressive cases and controls of the Radiant study sample.

	rs3746266A > G <sup>a</sup>			rs2075017T > C <sup>b</sup>		
	n	Estimates (95% C.I.)	p	n	Estimates (95% C.I.)	p
<b>All subjects<sup>c</sup></b>	2822	−0.67 (−1.13 to −0.21)	<b>0.004</b>	2136	−0.50 (−0.95 to −0.05)	<b>0.03</b>
Male	909	−0.38 (−1.05 to 0.30)	0.28	713	−0.25 (−0.93 to 0.42)	0.46
Female	1916	−0.79 (−1.38 to −0.21)	<b>0.008</b>	1424	−0.62 (−1.20 to −0.04)	<b>0.04</b>
<b>Subjects with MDD</b>	2138	−0.75 (−1.30 to −0.21)	<b>0.007</b>	1346	−0.59 (−1.20 to 0.02)	0.06
<b>Controls</b>	684	−0.41 (−1.20 to 0.39)	0.32	790	−0.32 (−0.95 to 0.31)	0.32

Model were adjusted for age and sex (when appropriate) and principal components C.I.: confidence interval, MDD: Major depressive disorder

<sup>a</sup> *CRTC1* rs3746266A > G and MDD status and/or sex interactions regarding BMI: p > 0.05.

<sup>b</sup> *CRTC1* rs2075017T > C and MDD status and/or sex interactions regarding BMI: p > 0.05.

<sup>c</sup> Available subjects with clinical and genetic data.

**Table 4**  
Association between *CRTC1* polymorphisms and BMI in the total, sex-stratified sample and in depressive cases and controls of the NESDA/NTR study sample.

	n	rs3746266A > G		rs6510997C > T	
		Estimates (SE)	p	Estimates (SE)	p
<b>All subjects</b>	4663	−0.047 (0.130)	0.73	0.014 (0.130)	0.91
Male	1676	0.072 (0.202)	0.72	−0.053 (0.193)	0.78
Female	2987	−0.121 (0.179)	0.50	0.046 (0.172)	0.79
<b>Subjects with MDD</b>	1768	0.038 (0.256)	0.88	0.166 (0.245)	0.50
<b>Controls</b>	2895	−0.076 (0.150)	0.61	−0.054 (0.144)	0.71

Model were adjusted for age and sex (when appropriate) and principal components. SE: standard error, MDD: Major depressive disorder.

and fat mass. However, the effect of the SNP was weaker compared to the psychiatric samples (Choong et al., 2013). In the present study, we were able to show in the psychiatrically evaluated sub-sample of CoLaus (PsyCoLaus) that the effect of *CRTC1* rs6510997C > T on fat mass was restricted to subjects with a lifetime history of MDD. In these subjects, the *CRTC1* rs6510997T-allele showed a significant protective effect for the fat mass. Although there was no significant interaction between the *CRTC1* SNPs and MDD, similar trend regarding the BMI was observed in the Radiant sample (no fat mass data available in this sample), which included more severe treated cases with lifetime recurrent depression. The *CRTC1* rs3746266A > G SNP only reached the level of statistical significance in MDD cases. However, these results could not be replicated in a third case-control sample with lifetime depression, the NESDA/NTR study.

*CRTC1* rs6510997C > T was associated with fat mass exclusively among depressive cases but not unaffected individuals in the PsyCoLaus sample. This association in depressed subjects did not differ in function of the depression subtypes (atypical versus others). Such an association with the BMI was not observed in the whole PsyCoLaus sample, but *CRTC1* SNPs were associated with BMI in the Radiant study. BMI may less accurately capture adiposity than estimated fat mass using bioimpedance (Marques-Vidal et al., 2009; Prentice and Jebb, 2001). Using the whole CoLaus sample, previous work established that fat mass enables capture of 3 times more subjects with high cardiovascular risk than BMI (Marques-Vidal et al., 2009) and we also showed that this sample is better powered to detect an association of the rs6510997C > T SNP with fat mass than with BMI (Choong et al., 2013). In the psychiatric samples, based on subjects at high risk of metabolic abnormalities because of the disorder and of the psychotropic medications, an association between *CRTC1* SNPs and BMI could be observed (Choong et al., 2013). Accordingly, the association

between *CRTC1* SNPs and BMI observed in the Radiant study could be explained by the higher degree of severity of the disorder in this sample compared to that in the PsyCoLaus. Interestingly, post hoc analysis also revealed a significant protective effect of the *CRTC1* rs6510997C > T SNP on BMI and waist circumference in PsyCoLaus subjects with MDD (but not in the overall sample) which would be in line with our hypothesis that the impact of *CRTC1* variants on adiposity markers is restricted to affected subjects.

*CRTC1* SNPs were associated with obesity markers in both PsyCoLaus and Radiant studies, but the association was attributable to MDD cases. On the other hand, *CRTC1* SNPs were not associated with BMI in the NESDA/NTR study, neither in the whole sample, nor in the stratified MDD case-control analyses. Previously, PsyCoLaus and DeCC (from the Radiant study) samples detected a protective effect of the *FTO* rs9939609-A variant on MDD (Samaan et al., 2013), while in the NESDA/NTR study the SNP was positively associated with MDD, especially in the atypical depression subtype (Milaneschi et al., 2014). Therefore, an inconsistency was already observed before for these studies which could be partly explained by the heterogeneity of depression in genetic association studies.

Obesity results from an imbalance between energy intake and energy expenditure. Recent studies have shown that mice lacking the *Crtc1* gene eat more and have less energy expenditure than wild-type mice, thus developing an obese feature, including obesity-related metabolic complications, under normal diets (Altarejos et al., 2008; Breuillaud et al., 2009). These results suggest that *CRTC1* is playing a major role in the hypothalamic control of food intake. *CRTC1* is mainly expressed in the brain (Altarejos et al., 2008; Breuillaud et al., 2009; Conkright et al., 2003; Kovacs et al., 2007; Wu et al., 2006) where it may modulate leptin anorexic effect in the hypothalamus. In the cell, the inactive phosphorylated form of *CRTC1* is sequestered in the cytoplasm, and its migration to the nucleus requires the concomitant activation of the phosphatase calcineurin and the inactivation of kinases of the 5' adenosine monophosphate-activated protein kinase (AMPK) family (Altarejos and Montminy, 2011). Interestingly, psychotropic drugs may increase weight by selective and potent stimulation of hypothalamic AMPK (Minokoshi et al., 2004) which has been shown to regulate food intake and reverse the actions of the anorexigenic hormone leptin (Kim et al., 2007).

In the PsyCoLaus sample, an interaction between *CRTC1* SNP and sex was observed regarding the fat mass and the protective effect of the SNP was mainly observed in females from the global sample. The same protective effect of *CRTC1* SNP on BMI was also observed in females from the global Radiant sample. This is in line with our previous results in the psychiatric and CoLaus samples (Choong et al., 2013). This stronger association found in females compared to males could be caused by a differential role of the leptin mediating satiety pathway in the enhancement of *CRTC1* activity. Women have much higher leptin levels than men (Rosenbaum et al., 1996) and female sex was found to predict stronger weight gain during psychotropic treatment (Gebhardt et al., 2009). A hypothetical mechanism of the effect of *CRTC1* and its interaction with sex hormones was published previously (Choong et al., 2013). Additionally, a meta-analysis of GWAS reported an association of an intronic SNP of *CRTC1* (rs10423674A > C) with the age of menarche (Elks et al., 2010), supporting a potential interaction between sex hormones and *CRTC1*.

The *CRTC1* SNPs were not associated with depression in our three MDD case-control studies. Animal models lacking the *Crtc1* gene showed behavioral abnormalities and depression-related behavior (Breuillaud et al., 2012), but this finding could not be translated in our study. However, the functional activities of these

SNPs in humans are still unknown and possibly the complete absence of the *CRTC1* gene could lead to the observed behavioral disturbances in animal models, which is probably not the case for effect of the tested SNPs on *CRTC1* expression. On the other hand, the observed effect of the *CRTC1* SNPs on obesity markers was uniquely detected in subjects with MDD. The lack of association between *CRTC1* SNPs and obesity markers in controls from the PsyCoLaus and the Radiant samples explains the weak association previously found in the population-based sample (Choong et al., 2013). It also confirms our hypothesis in which psychiatric illness and/or treatment with potentially weight gain-inducing psychotropic drugs could play a role in genetically mediated energy homeostasis and that the effect of *CRTC1* variations on obesity markers was unmasked in this group of subjects. Interestingly, a study in animal models (Mastronardi et al., 2011) showed that rats exposed to a high-fat diet after stress and treatment for a short period with antidepressants, to have more body weight and size from 17 to 22 weeks following antidepressant discontinuation compared to rats with the same conditions but treated with saline (Mastronardi et al., 2011). These findings support our concept of persistent, long-term effects of pharmacological–environment interactions on body weight regulation, even if the exposure to these medications was in any period in their lifetime.

This study has several limitations. Estimated fat mass using bioimpedance was only measured in the PsyCoLaus sample while only BMI was the only adiposity marker measured in the replication samples. Additionally, height and weight were self-reported in the Radiant study, which might influence the accuracy of BMI measurements in this sample, although the reliability of self-report of height and weight was assessed in the GENDEP dataset. Psychotropic medications were not available in the Radiant sample, therefore, associations between *CRTC1* SNPs and BMI in treated vs. untreated cases could not be calculated. Both *CRTC1* SNPs (rs6510997C > T and rs3746266A > G) were not always available in our three case-control samples; however, SNPs with nearly complete LD ( $r^2 > 0.9$ ) were selected and used as proxies, and results from these proxies could be generalized to our 2 *CRTC1* SNPs (meaning that the genotype present at one locus of the SNPs in LD is independent of the genotype at the second locus, both SNPs having the same allele frequency and could represent each other). On the other hand, SNPs were not always directly genotyped and different imputation methods with different imputation qualities were used which can potentially influence the results on the associations between different imputed *CRTC1* SNPs and obesity markers in our studies.

In conclusion, we showed in this study that *CRTC1* polymorphisms play no role in obesity markers in the general population and that the weak effect previously reported was driven by cases diagnosed with lifetime MDD. *CRTC1* seems to play an important role in the high prevalence of overweight and obesity specifically in psychiatric samples, a population at risk of developing obesity because of the disease itself and/or the medications. However, *CRTC1* seems not to be implicated directly in the development of psychiatric diseases.

#### Conflict of interest

CBE received honoraria for conferences or teaching CME courses from Astra Zeneca, Janssen-Cilag, Lundbeck, Merck Sharp & Dohme, Mepha, Otsuka, Servier and Vifor-Pharma in the past 3 years. MP was part of an advisory board of Lundbeck in the past 3 years.

All authors declare no conflict of interest in relation to the content of the paper.

## Author disclosure:

CBE received research support the Swiss National Research Foundation (320030-120686 and 324730-144064), from Takeda and from the Roche Organ Transplantation Research Foundation in the past 3 years. The CoLaus/PsyCoLaus study received financial contributions from GlaxoSmithKline, the Faculty of Biology and Medicine of Lausanne, the Swiss National Science Foundation (105993, 118308, 122661, 139468, 148401).

The Radiant study was funded by the Medical Research Council, UK. GlaxoSmithKline (G0701420) funded the DeNT study and they were co-funder with the Medical Research Centre for the GWAS of the whole sample. The GENDEP study was funded by a European Commission Framework 6 grant, EC Contract Ref.: LSHB-CT-2003-503428. This study presents independent research [part-] funded by the National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

Netherlands Study of Depression and Anxiety and Netherland Twin Register: funding was obtained from the Netherlands Organization for Scientific Research (NWO) and MagW/ZonMW grants Middelgroot-911-09-032, Spinozapremie 56-464-14192, Geestkracht program of the Netherlands Organization for Health Research and Development (ZonMW 10-000-1002), Center for Medical Systems Biology (CSMB, NOW Genomics), Genetic influences on stability and change in psychopathology from childhood to young adulthood (ZonMW 912-10-020), NBIC/BioAssist/RK (2008.024), Biobanking and Biomolecular Resources Research Infrastructure (BBMRI-NL, 184.021.007), VU University's Institute for Health and Care Research (EMGO+) and Neuroscience Campus Amsterdam (NCA); the European Science Council (ERC Advanced, 230374). Part of the genotyping and analyses were funded by the Genetic Association Information Network (GAIN) of the Foundation for the National Institutes of Health, Rutgers University Cell and DNA Repository (NIMH U24 MH068457-06), the Avera Institute, Sioux Falls, South Dakota (USA) and the National Institutes of Health (NIH R01 HD042157-01A1, MH081802, Grand Opportunity grants 1RC2 MH089951 and 1RC2 MH089995). Computing was supported by BiG Grid, the Dutch e-Science Grid, which is financially supported by NWO.

## Author Contributions:

Drs Preisig (PsyCoLaus), Rivera (Radiant) and Milaneschi (NESDA/NTR) had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

*Study concept and design:* Preisig and Eap.

*Acquisition of data:* Preisig, Rivera, Milaneschi, Willemsen

*Analysis and interpretation of data:* Quteineh, Eap, Preisig, Castelao, Gholam-Rezaee, Rivera, Milaneschi

*Drafting of the manuscript:* Quteineh

*Critical revision of the manuscript for important intellectual content:* Quteineh, Preisig, Rivera, Milaneschi, Castelao, Gholam-Rezaee, Vandenberghe, Saigi-Morgui, Delacrétaz, Cardinaux, Willemsen, Boomsma, Penninx, Ching-López, Conus, Eap

*Statistical analysis:* Castelao, Gholam-Rezaee, Rivera, Ching-López, Milaneschi

*Obtained funding:* Preisig, Eap

*Administrative, technical, and material support:* Preisig, Rivera, Milaneschi

*Study supervision:* Preisig and Eap.

## Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.jad.2016.03.031>.

## References

- WHO WMH-CIDI, 2015. The World Health Organization World Mental Health Composite International Diagnostic Interview.
- Afari, N., Noonan, C., Goldberg, J., Roy-Byrne, P., Schur, E., Golnari, G., Buchwald, D., 2010. Depression and obesity: do shared genes explain the relationship? *Depress Anxiety* 27, 799–806.
- Altarejos, J.Y., Goebel, N., Conkright, M.D., Inoue, H., Xie, J., Arias, C.M., Sawchenko, P.E., Montminy, M., 2008. The Creb1 coactivator Crtc1 is required for energy balance and fertility. *Nat. Med.* 14, 1112–1117.
- Altarejos, J.Y., Montminy, M., 2011. CREB and the CRC co-activators: sensors for hormonal and metabolic signals. *Nat. Rev. Mol. Cell Biol.* 12, 141–151.
- Blaine, B., 2008. Does depression cause obesity?: A meta-analysis of longitudinal studies of depression and weight control. *J. Health Psychol.* 13, 1190–1197.
- Boomsma, D.I., de Geus, E.J., Vink, J.M., Stubbe, J.H., Distel, M.A., Hottenga, J.J., Posthuma, D., van Beijsterveldt, T.C., Hudziak, J.J., Bartels, M., Willemsen, G., 2006. Netherlands Twin Register: from twins to twin families. *Twin Res. Hum. Genet.* 9, 849–857.
- Breuilaud, L., Halfon, O., Magistretti, P.J., Pralong, F.P., Cardinaux, J.R., 2009. Mouse fertility is not dependent on the CREB coactivator Crtc1. *Nat. Med.* 15, 989–990.
- Breuilaud, L., Rossetti, C., Meylan, E.M., Merinat, C., Halfon, O., Magistretti, P.J., Cardinaux, J.R., 2012. Deletion of CREB-regulated transcription coactivator 1 induces pathological aggression, depression-related behaviors, and neuroplasticity genes dysregulation in mice. *Biol. Psychiatry* 72, 528–536.
- Choong, E., Quteineh, L., Cardinaux, J.R., Gholam-Rezaee, M., Vandenberghe, F., Dobrin, M., Bondolfi, G., Etter, M., Holzer, L., Magistretti, P., Von Gunten, A., Preisig, M., Vollenweider, P., Consortium, T.G., team, T.O., Beckmann, J.S., Pralong, F.P., Waerber, G., Kutalik, Z., Conus, P., Bochud, M., Eap, C.B., 2013. Influence of CRT1 polymorphisms on body mass index and fat mass in psychiatric patients and in the general adult population. *Jama Psychiatry* 70, 1011–1019.
- Cohen-Woods, S., Gaysina, D., Craddock, N., Farmer, A., Gray, J., Gunasinghe, C., Hoda, F., Jones, L., Knight, J., Korszun, A., Owen, M.J., Sterne, A., Craig, I.W., McGuffin, P., 2009. Depression Case Control (DeCC) Study fails to support involvement of the muscarinic acetylcholine receptor M2 (CHRM2) gene in recurrent major depressive disorder. *Hum. Mol. Genet.* 18, 1504–1509.
- Comings, D.E., Gade, R., MacMurray, J.P., Muhleman, D., Peters, W.R., 1996. Genetic variants of the human obesity (OB) gene: association with body mass index in young women, psychiatric symptoms, and interaction with the dopamine D2 receptor (DRD2) gene. *Mol. Psychiatry* 1, 325–335.
- Conkright, M.D., Canettieri, G., Sreanor, R., Guzman, E., Miraglia, L., Hogensch, J.B., Montminy, M., 2003. TORCs: transducers of regulated CREB activity. *Mol. Cell* 12, 413–423.
- Elks, C.E., Perry, J.R., Sulem, P., Chasman, D.I., Franceschini, N., He, C., Lunetta, K.L., Visser, J.A., Byrne, E.M., Cousminer, D.L., Gudbjartsson, D.F., Esko, T., Feenstra, B., Hottenga, J.J., Koller, D.L., Kutalik, Z., Lin, P., Mangino, M., Marongiu, M., McArdle, P.F., Smith, A.V., Stolk, L., van Wingerden, S.H., Zhao, J.H., Albrecht, E., Corre, T., Ingelsson, E., Hayward, C., Magnusson, P.K., Smith, E.N., Ulivi, S., Warrington, N.M., Zgaga, L., Alavere, H., Amin, N., Aspelund, T., Bandinelli, S., Barroso, I., Berenson, G.S., Bergmann, S., Blackburn, H., Boerwinkle, E., Buring, J.E., Busonero, F., Campbell, H., Chanock, S.J., Chen, W., Cornelis, M.C., Couper, D., Coviello, A.D., d'Adamo, P., De Faire, U., De Geus, E.J., Deloukas, P., Doring, A., Smith, G.D., Easton, D.F., Eiriksdottir, G., Emilsson, V., Eriksson, J., Ferrucci, L., Folsom, A.R., Foroufard, T., Garcia, M., Gasparini, P., Geller, F., Gieger, C., Gudnason, V., Hall, P., Hankinson, S.E., Ferrel, L., Heath, A.C., Hernandez, D.G., Hofman, A., Hu, F.B., Illig, T., Jarvelin, M.R., Johnson, A.D., Karasik, D., Khaw, K.T., Kiel, D.P., Kilpelainen, T.O., Kolcic, I., Kraft, P., Launer, L.J., Laven, J.S., Li, S., Liu, J., Levy, D., Martin, N.G., McArdle, W.L., Melbye, M., Mooser, V., Murray, J.C., Murray, S.S., Nalls, M.A., Navarro, P., Nelis, M., Ness, A.R., Northstone, K., Oostra, B.A., Peacock, M., Palmer, L.J., Palotie, A., Pare, G., Parker, A.N., Pedersen, N.L., Peltonen, L., Pennell, C.E., Pharoah, P., Polasek, O., Plump, A.S., Pouta, A., Porcu, E., Rafnar, T., Rice, J.P., Ring, S.M., Rivadeneira, F., Rudan, I., Sala, C., Salomaa, V., Sanna, S., Schlessinger, D., Schork, N.J., Scuteri, A., Segre, A.V., Shuldiner, A.R., Soranzo, N., Sovio, U., Srinivasan, S.R., Strachan, D.P., Tammesoo, M.L., Tikkanen, E., Toniolo, D., Tsui, K., Tryggvadottir, L., Tyrer, J., Uda, M., van Dam, R.M., van Meurs, J.B., Vollenweider, P., Waerber, G., Wareham, N.J., Waterworth, D.M., Weedon, M.N., Wichmann, H.E., Willemsen, G., Wilson, J.F., Wright, A.F., Young, L., Zhai, G., Zhuang, W.V., Bierut, L.J., Boomsma, D.I., Boyd, H.A., Crisponi, L., Demerath, E.W., van Duijn, C.M., Econs, M.J., Harris, T.B., Hunter, D.J., Loos, R.J., Metspalu, A., Montgomery, G.W., Ridker, P.M., Spector, T.D., Streeten, E.A., Stefansson, K., Thorsteinsdottir, U., Uitterlinden, A.G., Widen, E., Murabito, J.M., Ong, K.K., Murray, A., 2010. Thirty new loci for age at menarche identified by a meta-analysis of genome-wide association studies. *Nat. Genet.* 42, 1077–1085.
- Farmer, A., Breen, G., Brewster, S., Craddock, N., Gill, M., Korszun, A., Maier, W., Middleton, L., Mors, O., Owen, M., Perry, J., Preisig, M., Rietschel, M., Reich, T., Jones, L., Jones, I., McGuffin, P., 2004. The Depression Network (DeNT) Study: methodology and sociodemographic characteristics of the first 470 affected sibling pairs from a large multi-site linkage genetic study. *BMC Psychiatry* 4, 42.

- Farmer, A., Korszun, A., Owen, M.J., Craddock, N., Jones, L., Jones, I., Gray, J., Williamson, R.J., McGuffin, P., 2008. Medical disorders in people with recurrent depression. *Br. J. Psychiatry* 192, 351–355.
- Farmer, A., Korszun, A., Owen, M.J., Craddock, N., Jones, L., Jones, I., Gray, J., Williamson, R.J., McGuffin, P., 2008. Medical disorders in people with recurrent depression. *Br. J. Psychiatry* 192, 351–355.
- Forty, L., Ulanova, A., Jones, L., Jones, I., Gordon-Smith, K., Fraser, C., Farmer, A., McGuffin, P., Lewis, C.M., Hosang, G.M., Rivera, M., Craddock, N., 2014. Comorbid medical illness in bipolar disorder. *Br. J. Psychiatry* 205, 465–472.
- Frayling, T.M., Timpson, N.J., Weedon, M.N., Zeggini, E., Freathy, R.M., Lindgren, C.M., Perry, J.R., Elliott, K.S., Lango, H., Rayner, N.W., Shields, B., Harries, L.W., Barrett, J.C., Ellard, S., Groves, C.J., Knight, B., Patch, A.M., Ness, A.R., Ebrahim, S., Lawlor, D.A., Ring, S.M., Ben-Shlomo, Y., Jarvelin, M.R., Sovio, U., Bennett, A.J., Melzer, D., Ferrucci, L., Loos, R.J., Barroso, I., Wareham, N.J., Karpe, F., Owen, K.R., Cardon, L.R., Walker, M., Hitman, G.A., Palmer, C.N., Doney, A.S., Morris, A.D., Smith, G.D., Hattersley, A.T., McCarthy, M.I., 2007. A common variant in the FTO gene is associated with body mass index and predisposes to childhood and adult obesity. *Science* 316, 889–894.
- Gebhardt, S., Haberhausen, M., Heinzl-Gutenbrunner, M., Gebhardt, N., Remschmidt, H., Krieg, J.C., Hebebrand, J., Theisen, F.M., 2009. Antipsychotic-induced body weight gain: predictors and a systematic categorization of the long-term weight course. *J. Psychiatry Res.* 43, 620–626.
- Hung, C.F., Breen, G., Czamara, D., Corre, T., Wolf, C., Kloiber, S., Bergmann, S., Craddock, N., Gill, M., Holsboer, F., Jones, L., Jones, I., Korszun, A., Kutalik, Z., Lucae, S., Maier, W., Mors, O., Owen, M.J., Rice, J., Rietschel, M., Uher, R., Vollenweider, P., Waeber, G., Craig, I.W., Farmer, A.E., Lewis, C.M., Muller-Mysok, B., Preisig, M., McGuffin, P., Rivera, M., 2015. A genetic risk score combining 32 SNPs is associated with body mass index and improves obesity prediction in people with major depressive disorder. *BMC Med.* 13, 86.
- Hung, C.F., Rivera, M., Craddock, N., Owen, M.J., Gill, M., Korszun, A., Maier, W., Mors, O., Preisig, M., Rice, J.P., Rietschel, M., Jones, L., Middleton, L., Aitchison, K.J., Davis, O.S., Breen, G., Lewis, C., Farmer, A., McGuffin, P., 2014. Relationship between obesity and the risk of clinically significant depression: Mendelian randomisation study. *Br. J. Psychiatry* 205, 24–28.
- Kim, S.F., Huang, A.S., Snowman, A.M., Teuscher, C., Snyder, S.H., 2007. Antipsychotic drug-induced weight gain mediated by histamine H1 receptor-linked activation of hypothalamic AMP-kinase. *PNAS* 104, 3456–3459.
- Kovacs, K.A., Steullet, P., Steinmann, M., Do, K.Q., Magistretti, P.J., Halfon, O., Cardinaux, J.R., 2007. TORC1 is a calcium- and cAMP-sensitive coincidence detector involved in hippocampal long-term synaptic plasticity. *Proc. Natl. Acad. Sci. USA* 104, 4700–4705.
- Lewis, C.M., Ng, M.Y., Butler, A.W., Cohen-Woods, S., Uher, R., Piro, K., Weale, M.E., Schosser, A., Paredes, U.M., Rivera, M., Craddock, N., Owen, M.J., Jones, L., Jones, I., Korszun, A., Aitchison, K.J., Shi, J., Quinn, J.P., Mackenzie, A., Vollenweider, P., Waeber, G., Heath, S., Lathrop, M., Muglia, P., Barnes, M.R., Whittaker, J.C., Tozzi, F., Holsboer, F., Preisig, M., Farmer, A.E., Breen, G., Craig, I.W., McGuffin, P., 2010. Genome-wide association study of major recurrent depression in the UK population. *Am. J. Psychiatry* 167, 949–957.
- Luppino, F.S., de Wit, L.M., Bouvy, P.F., Stijnen, T., Cuijpers, P., Penninx, B.W., Zitman, F.G., 2010. Overweight, obesity, and depression: a systematic review and meta-analysis of longitudinal studies. *Arch. Gen. Psychiatry* 67, 220–229.
- Marques-Vidal, P., Bochud, M., Mooser, V., Paccaud, F., Waeber, G., Vollenweider, P., 2009. Obesity markers and estimated 10-year fatal cardiovascular risk in Switzerland. *Nutr. Metab. Cardiovasc Dis.* 19, 462–468.
- Mastrorandi, C., Paz-Filho, G.J., Valdez, E., Maestre-Mesa, J., Licinio, J., Wong, M.L., 2011. Long-term body weight outcomes of antidepressant-environment interactions. *Mol. Psychiatry* 16, 265–272.
- McGuffin, P., Katz, R., Aldrich, J., 1986. Past and present state examination: the assessment of lifetime ever' psychopathology. *Psychol. Med.* 16, 461–465.
- McGuffin, P., Knight, J., Breen, G., Brewster, S., Boyd, P.R., Craddock, N., Gill, M., Korszun, A., Maier, W., Middleton, L., Mors, O., Owen, M.J., Perry, J., Preisig, M., Reich, T., Rice, J., Rietschel, M., Jones, L., Sham, P., Farmer, A.E., 2005. Whole genome linkage scan of recurrent depressive disorder from the depression network study. *Hum. Mol. Genet.* 14, 3337–3345.
- Milaneschi, Y., Lamers, F., Mbarek, H., Hottenga, J.J., Boomsma, D.I., Penninx, B.W., 2014. The effect of FTO rs9939609 on major depression differs across MDD subtypes. *Mol. Psychiatry* 19, 960–962.
- Minokoshi, Y., Alquier, T., Furukawa, N., Kim, Y.B., Lee, A., Xue, B., Mu, J., Foulfelle, F., Ferre, P., Birnbaum, M.J., Stuck, B.J., Kahn, B.B., 2004. AMP-kinase regulates food intake by responding to hormonal and nutrient signals in the hypothalamus. *Nature* 428, 569–574.
- Nurnberger Jr., J.I., Blehar, M.C., Kaufmann, C.A., York-Cooler, C., Simpson, S.G., Harkavy-Friedman, J., Severe, J.B., Malaspina, D., Reich, T., 1994. Diagnostic interview for genetic studies. Rationale, unique features, and training. NIMH Genetics Initiative. *Arch. Gen. Psychiatry* 51, 849–859.
- Penninx, B.W., Beekman, A.T., Smit, J.H., Zitman, F.G., Nolen, W.A., Spinhoven, P., Cuijpers, P., De Jong, P.J., Van Marwijk, H.W., Assendelft, W.J., Van Der Meer, K., Verhaak, P., Wensing, M., De Graaf, R., Hoogendijk, W.J., Ormel, J., Van Dyck, R., Consortium, N.R., 2008. The Netherlands Study of Depression and Anxiety (NESDA): rationale, objectives and methods. *Int. J. Methods Psychiatr. Res.* 17, 121–140.
- Preisig, M., Fenton, B.T., Matthey, M.L., Berney, A., Ferrero, F., 1999. Diagnostic interview for genetic studies (DIGS): inter-rater and test-retest reliability of the French version. *Eur. Arch. Psychiatry Clin. Neurosci.* 249, 174–179.
- Preisig, M., Waeber, G., Vollenweider, P., Bovet, P., Rothen, S., Vandeldeur, C., Guex, P., Middleton, L., Waterworth, D., Mooser, V., Tozzi, F., Muglia, P., 2009. The PsychoLaus study: methodology and characteristics of the sample of a population-based survey on psychiatric disorders and their association with genetic and cardiovascular risk factors. *BMC Psychiatry* 9, 9.
- Prentice, A.M., Jebb, S.A., 2001. Beyond body mass index. *Obes. Rev.* 2, 141–147.
- Qi, L., Kang, K., Zhang, C., van Dam, R.M., Kraft, P., Hunter, D., Lee, C.H., Hu, F.B., 2008. Fat mass and obesity-associated (FTO) gene variant is associated with obesity: longitudinal analyses in two cohort studies and functional test. *Diabetes* 57, 3145–3151.
- Rivera, M., Cohen-Woods, S., Kapur, K., Breen, G., Ng, M.Y., Butler, A.W., Craddock, N., Gill, M., Korszun, A., Maier, W., Mors, O., Owen, M.J., Preisig, M., Bergmann, S., Tozzi, F., Rice, J., Rietschel, M., Rucker, J., Schosser, A., Aitchison, K.J., Uher, R., Craig, I.W., Lewis, C.M., Farmer, A.E., McGuffin, P., 2012. Depressive disorder moderates the effect of the FTO gene on body mass index. *Mol. Psychiatry* 17, 604–611.
- Rosenbaum, M., Nicolson, M., Hirsch, J., Heymsfield, S.B., Gallagher, D., Chu, F., Leibel, R.L., 1996. Effects of gender, body composition, and menopause on plasma concentrations of leptin. *J. Clin. Endocrinol. Metab.* 81, 3424–3427.
- Samaan, Z., Anand, S.S., Zhang, X., Desai, D., Rivera, M., Pare, G., Thabane, L., Xie, C., Gerstein, H., Engert, J.C., Craig, I., Cohen-Woods, S., Mohan, V., Diaz, R., Wang, X., Liu, L., Corre, T., Preisig, M., Kutalik, Z., Bergmann, S., Vollenweider, P., Waeber, G., Yusuf, S., Meyre, D., 2013. The protective effect of the obesity-associated rs9939609A variant in fat mass- and obesity-associated gene on depression. *Mol. Psychiatry* 18, 1281–1286.
- Scuteri, A., Sanna, S., Chen, W.M., Uda, M., Albai, G., Strait, J., Najjar, S., Nagaraja, R., Orru, M., Usala, G., Dei, M., Lai, S., Maschio, A., Busonero, F., Mulas, A., Ehret, G. B., Fink, A.A., Weder, A.B., Cooper, R.S., Galan, P., Chakravarti, A., Schlessinger, D., Cao, A., Lakatta, E., Abecasis, G.R., 2007. Genome-wide association scan shows genetic variants in the FTO gene are associated with obesity-related traits. *PLoS Genet.* 3, 1200–1210.
- Sham, P.C., Sterne, A., Purcell, S., Cherny, S., Webster, M., Rijdsdijk, F., Asherson, P., Ball, D., Craig, I., Eley, T., Goldberg, D., Gray, J., Mann, A., Owen, M., Plomin, R., 2000. GENESiS: creating a composite index of the vulnerability to anxiety and depression in a community-based sample of siblings. *Twin Res.* 3, 316–322.
- Uher, R., Huezio-Diaz, P., Perroud, N., Smith, R., Rietschel, M., Mors, O., Hauser, J., Maier, W., Kozel, D., Henigsberg, N., Barreto, M., Placentino, A., Dernovsek, M.Z., Schulze, T.G., Kalember, P., Zobel, A., Czerski, P.M., Larsen, E.R., Souery, D., Giovannini, C., Gray, J.M., Lewis, C.M., Farmer, A., Aitchison, K.J., McGuffin, P., Craig, I., 2009. Genetic predictors of response to antidepressants in the GENDEP project. *Pharm. J.* 9, 225–233.
- Willemsen, G., de Geus, E.J., Bartels, M., van Beijsterveldt, C.E., Brooks, A.I., Es-tourgie-van Burk, G.F., Fugman, D.A., Hoekstra, C., Hottenga, J.J., Kluff, K., Meijer, P., Montgomery, G.W., Rizzu, P., Sondervan, D., Smit, A.B., Spijker, S., Suchiman, H.E., Tischfield, J.A., Lehner, T., Slagboom, P.E., Boomsma, D.I., 2010. The Netherlands Twin Register biobank: a resource for genetic epidemiological studies. *Twin Res. Hum. Genet.* 13, 231–245.
- Wing, J.K., Babor, T., Brugha, T., Burke, J., Cooper, J.E., Giel, R., Jablenski, A., Regier, D., Sartorius, N., 1990. SCAN. Schedules for clinical assessment in neuropsychiatry. *Arch. Gen. Psychiatry* 47, 589–593.
- Wu, Z., Huang, X., Feng, Y., Handschin, C., Gullicksen, P.S., Bare, O., Labow, M., Spiegelman, B., Stevenson, S.C., 2006. Transducer of regulated CREB-binding proteins (TORCs) induce PGC-1 $\alpha$  transcription and mitochondrial biogenesis in muscle cells. *Proc. Natl. Acad. Sci. USA* 103, 14379–14384.